ResearchGate

See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/328631207

Impact of packaging material and storage condition on retention of provitamin A carotenoids and xanthophylls in yellow-seeded maize flour

Article *in* Functional Foods in Health and Disease · October 2018

DOI: 10.31989/ffhd.v8i10.535

CITATIONS
7

READS
966

5 authors, including:



Wasiu Awoyale
Kwara State University
102 PUBLICATIONS **877** CITATIONS
SEE PROFILE



Emmanuel Oladeji Alamu
Nice-Pak Products Inc.
166 PUBLICATIONS **1,588** CITATIONS
SEE PROFILE



Emmanuel Anyachukwu Irondi
Kwara State University
79 PUBLICATIONS **1,033** CITATIONS
SEE PROFILE



Busie Maziya-Dixon
International Institute of Tropical Agriculture
246 PUBLICATIONS **5,338** CITATIONS
SEE PROFILE

All content following this page was uploaded by Wasiu Awoyale on 31 October 2018.

The user has requested enhancement of the downloaded file.

Research Article Open Access

Impact of packaging material and storage condition on retention of provitamin A carotenoids and xanthophylls in yellow-seeded maize flour

Wasiu Awoyale [1, 2], Emmanuel O. Alamu[1], Emmanuel A. Irondi[3], Busie Maziya-Dixon [1*] and Abebe Menkir[1]

[1]International Institute of Tropical Agriculture (IITA), Grosvenor House, Croydon, CR09 XP, United Kingdom; [2]Department of Food Science & Technology, Kwara State University Malete, PMB 1530, Ilorin Kwara State, Nigeria; [3]Department of Biochemistry, Kwara State University, Malete, P.M.B. 1530, Ilorin, Nigeria

Corresponding author: Dr. Busie Maziya-Dixon. International Institute of Tropical Agriculture (IITA), Ltd. Grosvenor House, Croydon, CR09 XP, United Kingdom

Submission date: July 20th, 2018**, Acceptance Date:** October 28th, 2018, **Publication Date:** October 30th, 2018

Citation: Awoyale W., Alamu E.O., Irondi E.A., Maziya-Dixon B., Menkir A. Impact of packaging material and storage condition on retention of provitamin A carotenoids and xanthophylls in yellow-seeded maize flour. *Functional Foods in Health and Disease* 2018; 8(10): 462-477. https://doi.org/DOI#

ABSTRACT

Background: Pro-vitamin A carotenoid (pVAC) rich foods are those foods that contain substance which can be converted within the human body into retinol. These foods also contribute to the reduction of vitamin A deficiency diseases. Yellow-seeded maize flour is a pVAC rich food. Identifying the right packaging materials and storage conditions that retain pVAC in this food is essential for their health benefits. Traditionally, maize flour is stored with different packaging materials to increase its shelf life. For example, previous studies have shown how during storage in different food matrices, carotenoids, including maize grains, are highly susceptible to degradation by temperature, light, and oxygen. Therefore, in this study we investigated the effect of storage packaging materials (polypropylene woven sacks-PWS, high-density polyethylene bags-HDPE, and polyvinyl plastic containers-PPC) and storage condition (temperature and relative humidity) on retaining pVAC in yellow-seed maize flour.

Methods: The yellow-seeded maize grains were collected and ground into flour. The maize flour was divided into portions (200 g). Each portion was packed and sealed in PWS, HDPE, and PPC. The control samples (12 pieces) were stored on top of the storage box. The packed samples were stored in both the upper (12 samples) and lower (12 samples) compartment of a storage wooden box. The interior of the upper part was lightened with aluminum foil and fitted with fluorescent tube to increase the light intensity. The lower compartment was darkened with gloss black

painting. The flour samples were stored for 28 days. Samples were collected for pVAC and xanthophylls analyses at 7 day intervals using standard methods.

Results: The results demonstrated how packaging in PPC and storing in a dark compartment resulted in the highest total pVAC (92.39%) and total xanthophylls (89.44%) retention and retinol equivalent (RE) (0.40 μg/g). In contrast, packaging in HDPE and storing in lighted compartment resulted in the lowest pVAC (44.92%) and total xanthophylls (46.76%) retention and RE (0.19 μg/g).

Conclusions: Packaging yellow-seeded maize flour in PPC and storing in the dark may be recommended for maximum retention of carotenoids in yellow-seeded maize flour since, as the packaging material and storage condition resulted in the highest pVAC retention and retinol equivalent.

Keywords: Yellow-seeded maize; Carotenoids; Packaging materials; Storage conditions; Retinol equivalent

BACKGROUND

Carotenoids, the lipid-soluble plants secondary metabolites with 30 and 40 carbon atoms, are important in human nutrition and health because of their various benefits. Carotenoids are significant for their antioxidative properties, protecting cells against free radicals and reactive oxygen species [1], in addition to the oxidative damage they cause to cellular biomolecules such as deoxyribonucleic acid, proteins, and lipids [2]. Furthermore, carotenoids possess several other health benefits including provitamin A and immunostimulatory [3], anti-hepatocellular cancer [4], anti-inflammatory and antiapoptotic activities [5], and prevention of cataracts and age-related macular degeneration [6]. Carotenoids also contribute to organoleptic and nutritional properties of food and food by-products, providing the red, yellow, or orange coloring of fruits and vegetables. They are used as food colorants in foods and animal products such as egg yolk, butter, crustaceans, trout, salmon, and shrimp [7]. Through their structure, carotenoids are classified into two main groups: (i) carotenes also called carotenoid hydrocarbons, which only contain carbon and hydrogen, and (ii) xanthophylls or oxygenated carotenoids that may contain different functional groups (epoxy, methoxy, hydroxy, carbonyl and carboxyl acid groups) [8]. However, humans and other animals are unable to synthesize carotenoids. Consequently, they depend on the consumption of plant dietary products to meet their carotenoids demands for the various physiological and health benefits they provide [9].

Among the various cereal dietary sources of carotenoids, maize (Zea mays L.), especially the yellow-seeded type, is notable as the cereal that accumulates the highest level of carotenoids, having up to 80 μg total carotenoids per gram dry weight [10]. The yellow-seeded maize contains both provitamin A carotenoids (pVAC - β-carotene, α-carotene, and β-cryptoxanthin) and xanthophylls (lutein and zeaxanthin) without provitamin A activity [11]. However, the provitamin A content of maize varies, with yellow maize varieties has 0.5 to 1.5 μg/g provitamin A [12], which is insufficient to prevent vitamin A deficiency (VAD) in areas where diets are dominated by maize and maize products. For this reason, breeding programs aim to increase the level of

pVAC in yellow maize varieties from below 2 µg to 15 µg/g pVAC/g in orange grain varieties through biofortification [10]. However, the processing of the maize grain into floury products may reduce the quantity of the pro-vitamin A carotenoid (pVAC).

Traditionally, maize flour is stored in different packaging materials to increase its shelf life. The research carried out by De Moura et al [13] and Rodriguez-Amaya [14] revealed that carotenoids in stored maize grain foods are highly susceptible to degradation by temperature, light, and oxygen during storage. Bechoff et al. [15] and Alves et al. [16] further demonstrated that oxygen contributed more to the degradation of carotenoids than the temperature in food matrixes having high exposure to oxygen. Other studies also indicated that depending on the genotype when the grains of maize are stored for a period of six months using traditional methods, the degradation of pVAC in the grains may reach 60% [13, 17]. Additionally, previous studies reported 78–100% pVAC retention when cooking freshly ground biofortified orange maize meal to prepare Nshima [18, 19]. Stability of carotenoids during storage of biofortified maize has been studied by Mugode et al. [19]. These researchers reported that most of the carotenoid degradation occurred in the first weeks of storage, and then afterwards the degradation rate lowered. This makes it imperative to identify storage conditions and packaging materials that can maximally retain both pVAC and xanthophylls in flour of yellow-seeded maize during storage.

Therefore, this study was designed to investigate the impact of different packaging materials and storage conditions on the retention of pVAC and xanthophylls in yellow-seeded maize flour.

MATERIALS AND METHODS

Materials

Packaging materials

The various packaging materials, including polypropylene woven sacks (PWS), high-density polyethylene bags (HDPE), and polyvinyl plastic containers (PPC) were produced by Afriplast Industries Ltd, Oke-Bola, Ibadan, Nigeria. The properties of the packaging materials as specified by the manufacturer (Afriplast Industries Ltd, Sw7/8, Obafemi Awolowo Way, Oke-Bola, Ibadan, Nigeria) are shown in Table 1 [20].

Table 1. Properties of the packaging materials

Packaging material	Thickness (µm)	Oxygen permeability at 25 °C/24 h (mm/100 cm^2)	Water vapor permeability at 37.8 °C/24 h (mm/100 cm^2)	Relative humidity (%)
PWS	0.75	160.00	0.27	90.00
HDPE	1.30	500.00	1.40	90.00
PPC	0.45	80.00	8.00	90.00

PWS-polypropylene woven sacks, HDPE-high-density polyethylene bags, PPC-polyvinyl plastic containerS

Chemicals and reagents

Chemicals, reagents, and solvents used for the carotenoids profiling were all HPLC grade. Carotenoids standards (9-cis-β-carotene, 13-cis-β-carotene, all-trans-β-carotene, β-cryptoxanthin, lutein, and zeaxanthin) were bought from CaroteNature, GmbH (Lupsingen, Switzerland). All other chemicals, reagents, and solvents were of analytical grade.

Sample collection and preparation

A sample of yellow maize (DMR-LSRY) grains were collected from the Maize Improvement Programme of the International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. The grains were sorted and ground into flour using a laboratory hammer grinding machine fitted with a 0.5 mm mesh sieve. The flour was immediately used for the storage study.

Storage study

The storage study was conducted in a cubic wooden box of 2.6 ft length, width, and height, with upper and lower compartments. The interior part of the upper compartment was lined with aluminum foil and fitted with a T8 LongLast™ fluorescent tube (2 ft, 18 W, & 3000 K warm white) to increase light intensity. Gloss black paint was used to paint the lower compartment to maintain a dark enclosure for the samples. Each compartment had a separate door, for the ease of sample collection. The packaging materials, specifically the HDPE and PWS bags, were prepared to be used for packaging by cutting them into required sizes using a scissor and shaped using electric sealing and stitching machines respectively. The sizes of the packaging materials are: PWS (25 cm height × 13 cm breadth) and HDPE (23 cm height × 16 cm breadth).

The PPC size was measured to be 6 cm height ×13 cm breadth. A portion (200 g) of the flour was properly weighed using a weighing scale, packed, and sealed in each of the three different packaging materials as the following: PWS was sealed with a stitching machine, HDPE was sealed with an electric sealer, and PPC was covered hermetically with a lid [21]. The packaged flour samples replicated four times were stored in both the upper (lighted) and lower (dark) compartments of the storage box. The control flour samples also replicated four times were stored on top of the storage box. The flour samples (36 pieces) were stored for 28 days, during which samples were collected from each package at 7 day intervals for carotenoids profiling. The ambient temperature and relative humidity of each storage condition were measured with thermo-hygrometer (Max-Min) prior to each batch of sample collection (Table 2).

Table 2. Temperature and Relative humidity of each of the storage compartment

Compartment	3 weeks		6 weeks		9 weeks		12 weeks	
	Temp.(^{o}C)	R.H (%)	Temp.(^{o}C)	R.H (%)	Temp.(^{o}C)	R.H (%)	Temp.(^{o}C)	R.H (%)
Light	33.7	54.0	33.8	50.0	34.2	52.0	35.2	50.0
Dark	27.4	66.0	27.2	58.0	28.0	63.0	30.0	58.0
Outside box	26.6	75.0	26.1	63.0	27.5	68.0	27.9	73.0

Temp- Temperature, R.H-Relative humidity

Methods

Analysis of provitamin A carotenoids (pVAC) and xanthophylls using HPLC-PDAD

The pVAC and xanthophylls profiles of the flour samples were quantified according to the method described by Howe and Tanumihardjo [22], using high performance liquid chromatography system coupled with photodiode array detector (HPLC-PDAD). The HPLC instrumentation comprised Waters HPLC system (Water Corporation, Milford, MA), binary HPLC pump (Waters 626), auto-sampler (Waters 717), PDAD (Waters 2996) and Empower 1 software (Waters Corporation). Carotenoids were extracted from the flours by mixing 0.6 g of sample with 10 mL of ethanol containing 0.1% butylated hydroxyl toluene (BHT) and allowing the mixture to stand in a water bath at 85 °C for 5 min [22]. Subsequently, interfering oil in the mixture was saponified by adding 500 µL of potassium hydroxide (80% w/v). Afterward, the mixture was returned to the water bath (at 85 °C for 5 min) for two successive cycles, with thorough mixing after each cycle.

Then, the samples were placed in an ice bath and cold deionized water (3 mL) was added. The carotenoids in the samples mixture were separated for three consecutive times by adding 3 mL of hexane and then centrifuging at 1200 g for 5 min. The collected hexane fraction was washed three times with deionized water, centrifuged at 1200 g for 5 min, and dried under nitrogen gas using TurboVap LIV concentrator. Subsequently, the dried extract was re-dissolved in 1 mL methanol/dichloromethane (50:50 v/v), out of which an aliquot of 100 µL was injected into the HPLC system for the quantification of the carotenoids.

The mobile phase consisted of solvent A [methanol: water (92:8 v/v), with 10 mmol/L ammonium acetate] and solvent B (100% methyl tertiary-butyl ether). Gradient elution programme, set at 1 mL/min, was as follows: linear gradient from 83% to 59% A for 29 min; linear gradient from 59% to 30% A for 6 min; 1 min hold at 30% A; linear gradient from 30% to 83% A for 4 min; and 4 min hold at 83%. Chromatographic separation of carotenoids was carried out using a C30 YMC guard column (4.6 9 250 mm, 3 µm). Identifications and quantifications of carotenoids were done at 450 nm using an external standard method by comparing their retention times and spectra with those of reference standards. Samples were analyzed in triplicate.

Estimation of total provitamin A carotenoid content

Total pVAC was calculated using the formula reported by Taleon et al. [23]:

Total pVAC = βC + (1/2) (13-cis-βC) + (1/2) (9-cis-βC) + (1/2) (βCX);

βC is β-carotene, 13-cis-βC is 13-cis-β-carotene, 9-cis-βC is 9-cis-β-carotene, and βCX is β-cryptoxanthin.

Retention of provitamin A carotenoid

The percentage apparent retention (%AR) of total pVAC of the samples was calculated using the formula reported by Li et al. [24] as the following:

%AR = (pVAC per g of maize after storage x 100) / (pVAC per g of maize before storage)

Calculation of retinol equivalent of the provitamin A carotenoid

Retinol equivalent (RE) of the pVAC, expressed in μg per g dry weight, was calculated according to the in-vivo conversion factor proposed by the world health organization and reported by Djuikwo et al. [25], where 6 μg of β-carotene and 12 μg of α-carotene correspond to 1 RE. The RE of 13-cis-β-carotene and 9-cis-β-carotene was computed at the same strength equivalence as β-carotene.

Statistical analysis

Result data, expressed as the mean ± standard deviation, were analyzed statistically using analysis of variance (ANOVA) and least significant difference tests using SPSS statistical software package (version 17) at 95% confidence level.

RESULTS AND DISCUSSION

Carotenoids are unstable compounds. Due to their polyene structure, they act as antioxidants, a property linked to their own destruction involving abiotic factors such as temperature, light, metal catalysts, and water content. However, the same structural attributes of carotenoids which make them beneficial to human health also make them subject to oxidation. Oxidizing environments leading to non-enzymatic cleavage can also be initiated by cellular activities. Co-oxidation of carotenoids as a consequence of lipoxygenase activities is a long-known example of the non-enzymatic cleavage of carotenoids, in which the fatty acyl peroxy radicals generated cleave carotenoids by a random attack of the polyene chromophore [26, 27].

In this study, the pVAC (β-cryptoxanthin; α-carotene; 13-cis-β-carotene; 9-cis-β-carotene, all-trans-β-carotene, total β-carotene and total pVAC) profiles (μg/g dry weight) of the yellow-seeded maize grain flour as affected by storage period is presented in Table 3. The result shows that β-cryptoxanthin and all-trans-β-carotene were the predominant pVAC in the flour before and on the 28th day of storage (2.22, 1.63 μg/g and 0.77, 0.56 μg/g respectively). Generally, the levels of all the pVAC decreased as the storage period increased. However, while the decrease of the carotenoids during storage was significant ($p < 0.05$) for α-carotene, all-trans-β-carotene, 9-cis-β-carotene, total β-carotene and total pVAC, for β-cryptoxanthin and 13-cis-β-carotene it was not significant ($p > 0.05$). Similarly, the various packaging materials influenced the pVAC profile of the flour during the storage period. The effect of the packaging materials on the carotenoids was significant ($p<0.05$) for β-cryptoxanthin, 13-cis-β-carotene, all-trans-β-carotene, total β-carotene, the total pVAC and α-carotene, but not significant ($p > 0.05$) for 9-cis-β-carotene. Overall, storage period and packaging material combined only had a significant ($p<0.05$) effect on the level of all-trans-β-carotene (Table 3).

The range of the various pVAC observed in this study agrees with the values reported by other previous studies for the pVAC content of grains of yellow maize varieties [19, 17, 23]. The preponderance of β-cryptoxanthin and all-trans-β-carotene over other pVAC both before and during storage is also in agreement with an earlier report by Mugode et al. [19], indicating that these two carotenoids are the major pVAC in dry yellow maize grains, even with the use of different packaging materials and storage conditions in the present study. Previous studies have reported a decrease in the pVAC content of the yellow maize grain during storage [19, 23], which is attributed to exposure to high temperature, oxygen, and light that cause carotenoids degradation

during storage [13, 14]. Recently, Awoyale et al. [20] also reported a decrease in the pVAC content of Ogi powder made from yellow maize grain during storage. Therefore, the decrease in the pVAC content of the yellow maize flour with increasing storage period observed in this study is inconsonant with the existing literature on the behavior of pVAC in both yellow maize grain and its products during storage, at different temperatures and relative humidity. Additionally, it is important to note the measurement of the temperatures and relative humidity of the storage box was used to check the effect of lighting and darkness on the storage condition/atmosphere of the samples vis-à-vis carotenoid degradation.

Table 3. Provitamin A carotenoids (pVAC) profile (µg/g dry weight) of yellow maize grain flour as affected by storage period and packaging material

pVAC	Storage period (days)	Average (µg/g)	Range (µg/g)	P Storage period	P Packages	P Storage Period x Packages
βCX	0	2.22				
	7	1.95	1.52 - 2.74	NS	**	NS
	14	1.70	1.34 - 2.02	NS	**	NS
	21	1.71	1.43 - 2.04	NS	**	NS
	28	1.63	0.94 - 2.51	NS	**	NS
αC	0	0.34				
	7	0.30	0.22 - 0.43	*	***	NS
	14	0.25	0.19 - 0.29	*	***	NS
	21	0.23	0.17 - 0.29	*	***	NS
	28	0.16	0.03 - 0.27	*	***	NS
13-cis-βC	0	0.17				
	7	0.17	0.12 - 0.21	NS	**	NS
	14	0.15	0.10 - 0.17	NS	**	NS
	21	0.13	0.09 - 0.21	NS	**	NS
	28	0.11	0.06 - 0.27	NS	**	NS
9-cis-βC	0	0.35				
	7	0.30	0.08 – 0.44	**	NS	NS
	14	0.28	0.22 – 0.33	**	NS	NS
	21	0.28	0.23 – 0.34	**	NS	NS
	28	0.27	0.15 – 0.39	**	NS	NS
trans-βC	0	0.77				

	7	0.66	0.51 – 0.79	*	**	*
	14	0.58	0.46 – 0.66	*	**	*
	21	0.60	0.47 – 0.70	*	**	*
	28	0.56	0.33 – 0.90	*	**	*
Total βC	0	1.29				
	7	1.12	0.89 – 1.41	**	**	NS
	14	1.00	0.80 - 1.17	**	**	NS
	21	1.01	0.80 – 1.25	**	**	NS
	28	0.94	0.53 – 1.56	**	**	NS
Total pVAC	0	2.57				
	7	2.24	1.76 - 2.96	**	**	NS
	14	1.98	1.59 - 2.26	**	**	NS
	21	1.98	1.61 - 2.42	**	**	NS
	28	1.84	1.04 - 2.95	**	**	NS

Data represent the results of triplicate analyses. *$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$; NS - not significant ($p > 0.05$); P- p-value; βCX - β-cryptoxanthin; αC – α-carotene; 13-*cis*-βC - 13-*cis*-β-carotene; 9-*cis*-βC - 9-*cis*-β-carotene; *trans*-βC – all-*trans*-β-carotene; βC – β-carotene; pVAC - provitamin A carotenoids.

The xanthophylls (lutein, zeaxanthin, and total xanthophylls) profiles (μg/g dry weight) of the yellow maize grain flour as affected by storage period is presented in Table 4. As with the pVAC, the levels of the xanthophylls decreased with increasing storage period. Lutein decreased from 5.76 μg/g to 3.75 μg/g; zeaxanthin decreased from 10.61 μg/g to 8.18 μg/g; the total xanthophylls decreased from 16.37 μg/g to 11.94 μg/g, before storage (0 days) and at the 28th day of storage respectively. However, while the decrease was significant ($p<0.05$) for lutein and the total xanthophylls, it was not significant ($p > 0.05$) for zeaxanthin. The various packaging materials had a significant ($p<0.05$) effect on the levels of lutein, zeaxanthin, and total xanthophylls. Altogether, storage period and packaging material had significant ($p< 0.05$) effect on the levels of lutein and the total xanthophylls but no significant ($p > 0.05$) effect on the level of zeaxanthin. The levels of xanthophylls (lutein and zeaxanthin) quantified in the maize flours packaged in different materials and stored for 28th days under different conditions (Table 4) were consistently higher than the levels of their counterpart pVAC (Table 3). This observation is consistent with some previous reports that indicated that xanthophylls are the major carotenoids in yellow maize varieties [17, 23]. However, among the various carotenoids (both pVAC and xanthophylls), zeaxanthin was the most abundant carotenoid in the flours. This observation is in conformity with an earlier report by

Ortiz et al. [17], who reported that zeaxanthin was the predominant carotenoid in the different maize genotypes they evaluated in their study. As noted for the pVAC, the decrease in the xanthophylls may also be attributed to carotenoids degradation during storage because of exposure to high temperature, oxygen, and light [13, 14].


Table 4. Xanthophylls profile (µg/g dry weight) of yellow maize grain flour as affected by storage period and packaging materials

Xanthophylls	Storage period (days)	Average (µg/g)	Range (µg/g)	P Storage period	P Packages	P Storage period x Packages
Lut	0	5.76				
	7	5.08	4.19 – 5.99	***	**	*
	14	4.70	3.78 - 5.59	***	**	*
	21	4.18	3.23 - 4.71	***	**	*
	28	3.75	2.08 - 5.42	***	**	*
Zeax	0	10.61				
	7	8.78	7.41 - 10.56	NS	**	NS
	14	8.12	6.34 - 9.77	NS	**	NS
	21	7.68	1.67 - 9.77	NS	**	NS
	28	8.18	4.93 - 12.91	NS	**	NS
Total xanto	0	16.37				
	7	13.64	11.65 - 16.55	*	**	*
	14	12.82	10.12 - 15.29	*	**	*
	21	11.85	6.27 - 14.47	*	**	*
	28	11.94	7.01 - 18.01	*	**	*

Data represent the results of triplicate analyses. *$p \leq 0.05$; **$p \leq 0.01$; ***$p \leq 0.001$; NS - not significant ($p > 0.05$); P- p-value; Lut - lutein; Zeax - zeaxanthin; Xanto – xanthophylls.

The percentage apparent retention (%AR) of total pVAC in the yellow maize grain flour as affected by packaging materials, storage period, and storage conditions is depicted in Figure 1. At the end of the 28 day storage period, both packaging materials and storage period had a significant effect ($p < 0.05$) on the retention of the total pVAC of the maize flour. Generally, storage in the dark compartment of the storage box resulted in higher retention of total pVAC, when compared to storage in the lighted compartment of the box and storage outside the box, regardless of the packing material used. Furthermore, packaging in PPC led to higher retention of the pVAC relative to HDPE and PWS, regardless of the storage period and condition. Maize flour packaged in PPC and stored in dark compartment of the storage box (MCD) had the highest ($p < 0.05$) retention of total pVAC (92.39%) relative to the other packaging materials and storage conditions. In contrast, maize flour packaged in HDPE and stored in a lighted compartment of the storage box (MNL) had the least total pVAC retention (44.92%).

Figure 1. True retention (%TR) of total provitamin A carotenoids (pVAC) in yellow maize grain flour as affected by packaging materials, storage period and storage condition




The trend of the total pVAC retention as the storage period increased was not definite. While there was a steady decrease in the retention of the total pVAC for the various packaging materials and storage conditions from the zero (0) day to the 14th day of the storage, the decrease remained progressive for packaging in PWS and HDPE until the 28th day, regardless of the storage condition. However, packaging in the PPC resulted in an increase in the total pVAC retention on the 28th day for storage both in the dark compartment of the box and storage outside the box.

Degradation of pVAC, leading to their low retention in yellow maize grain [23] and its product (Ogi) [20] during storage has been reported before. The differential retention of total pVAC during the storage period by different packaging materials under different storage conditions could be explained by the nature of the packaging materials in relation to temperature, oxygen, and light. For the maize flour packaged in PPC and stored in dark compartment of the storage box, which had the highest pVAC retention, it is possible that the PPC and the dark compartment of the storage box provided an environment that excluded light and reduced exposure to heat. On the contrary, the lowest retention of pVAC in maize flour packaged in HDPE and stored in a lighted compartment of the storage box may be attributed to light and heat (ambient temperature of 35 °C) in that compartment [20, 28].

The percentage apparent retention (%AR) of total xanthophylls in yellow maize grain flour as affected by packaging materials, storage period and storage condition is shown in Figure 2. The Figure shows that total xanthophylls retention varied, depending on the storage period, packaging material and storage condition. At the end of the 28 day storage period, maize flour packaged in PPC and stored in dark compartment of the storage box (MCD) had the highest ($p < 0.05$) retention of total xanthophylls (89.44%), followed by maize flour packaged in PPC and stored outside the box (MCO) (85.85%); maize flour packaged in HDPE and stored in lighted compartment of the

storage box (MNL) had the least total xanthophylls (46.76%). Total xanthophylls retention decreased progressively from the zero (0) day to the 28th day of storage in maize flour packaged in HDPE and stored in both the lighted compartment (MNL) and the dark compartments (MND) of the storage box, with maize flour packaged in PWS and stored in dark compartment (MNL).

Figure 2. True retention (%TR) of total xanthophylls in yellow maize grain flour as affected by packaging materials, storage period and storage condition



MNO - maize flour packaged in high-density polyethylene bag and stored outside the box;
MSO - maize flour packaged in polypropylene woven sack and stored outside the box;
MCO - maize flour packaged in polyvinyl chloride container and stored outside the box;
MND - maize flour packaged in high-density polyethylene bag and stored in the dark compartment;
MSD - maize flour packaged in polypropylene woven sack and stored in the dark compartment;
MNL - maize flour packaged in high-density polyethylene bag and stored in the lighted compartment;
MSL - maize flour packaged in polypropylene woven sack and stored in the lighted compartment;
MCD - maize flour packaged in polyvinyl chloride container and stored in the dark compartment;
MCL - maize flour packaged in polyvinyl chloride container and stored in the lighted compartment.

In contrast, total xanthophylls retention decreased steadily until the 21st day of the storage but increased at the 28th day in maize flour packaged in PPC and stored in dark compartment of the

storage box (MCD) and outside the storage box (MCO), in addition to within maize flour packaged in HDPE and stored outside the box (MNO). As explained for the pVAC, the higher retention of total xanthophylls in the maize flour packaged in PPC and stored in dark compartment of the storage box may be due to the exclusion of light and heat. However, this was not the case for the maize flour packaged in HDPE and stored in the lighted compartment of the storage box, which had the least retention. A comparison of the retention of the total pVAC (Figure 1) and of the total xanthophylls (Figure 2) indicates that regardless of the storage period, packaging material, and storage condition, there was a higher retention of total pVAC (92.39%) than total xanthophylls (89.44%).


This may be due to the unsaturated (double) bonds present in the carbon chain of the xanthophylls structure, which makes them more susceptible to degradation during processing and storage [29-31]. In fact, oxidative degradation, which is dependent on factors including the presence of oxygen, light, heat, enzymes, and co-oxidation with lipid hydroperoxides and metals [32, 33] has been identified as the principal cause of more extensive degradation of carotenoids. It was reported by Otten et al. [34] that the biological value of food materials with vitamin A activity is expressed as retinol equivalent (RE). Figure 3 shows the RE (μg/g) of total pVAC in yellow maize grain flour as influenced by packaging materials, storage period, and storage condition. Maize flour (not subjected to packaging and storage) had the highest RE (0.43 μg/g). The RE decreased as the storage period increased, regardless of the packaging material and storage condition.

However, the extent of the decrease varied depending on the packaging material and storage condition. As expected, among the various packaging materials and storage conditions, MCD (with the highest total pVAC retention) had the highest RE (0.40 μg/g), while MNL (with the least total pVAC retention) had the lowest RE (0.19 μg/g) (Figure 1). Generally, at the 28^{th} day of storage, maize flour packaged in PPC displayed the highest RE in the three different storage conditions used in the study.

As stated earlier, due to the unsaturated (double) bonds present in the carbon chain of carotenoids, the presence of light, heat, and oxygen are known to precipitate some chemical reactions such as isomerization (cis to trans) and oxidation during processing and storage of foods, leading to their degradation and biological activity reduction [29, 31]. This could explain why the maize flour packaged in PPC and stored in dark compartment of the storage box (MCD) had the highest RE (Figure 3), as there might have been retardation in the rate of oxidation and isomerization of the pVAC in the flour, due to the exclusion of light and reduction in exposure to heat.

In contrast, that maize flour packaged in HDPE and stored in lighted compartment of the storage box (MNL) had the lowest RE could indicate that both the light and the heat (ambient temperature of 35 °C) in that compartment may have enhanced the oxidation and isomerization of the pVAC in the flour, causing their degradation and loss of biological activity, and, in this case, retinol activity.

Figure 3. Retinol equivalent (µg/g) of total pVAC in yellow maize grain flour as affected by packaging materials and storage period



MNO - maize flour packaged in high-density polyethylene bag and stored outside the box;
MSO - maize flour packaged in polypropylene woven sack and stored outside the box;
MCO - maize flour packaged in polyvinyl chloride container and stored outside the box;
MND - maize flour packaged in high-density polyethylene bag and stored in the dark compartment;
MSD - maize flour packaged in polypropylene woven sack and stored in the dark compartment;
MNL - maize flour packaged in high-density polyethylene bag and stored in the lighted compartment;
MSL - maize flour packaged in polypropylene woven sack and stored in the lighted compartment;
MCD - maize flour packaged in polyvinyl chloride container and stored in the dark compartment;
MCL - maize flour packaged in polyvinyl chloride container and stored in the lighted compartment.

CONCLUSIONS

Xanthophylls were more abundant than provitamin A carotenoids (pVAC). Among the pVAC, β-cryptoxanthin and all-trans-β-carotene were predominant in the biofortified yellow maize flour, both before and during storage. Packaging materials and storage periods affected the retention of both the pVAC and xanthophylls and the retinol equivalent (RE) of the pVAC during the storage period. Packaging in polyvinyl chloride container and storing in dark compartment of the storage box resulted in the highest total pVAC and total xanthophylls retention and RE; in contrast, packaging in high-density polyethylene bag and storing in a lighted compartment of the storage box resulted in the least total pVAC and total xanthophylls retention, and RE. Thus, packing in polyvinyl chloride container and storing in the dark is recommended for maximum retention of

pVAC, xanthophylls, and retinol activity in yellow maize flour to ensure its optimum health benefits.

Competing Interests: The authors declare no conflict of interest.

Authors' Contributions: WA, BMD and AM designed the research, WA, EOA, BMD and AM performed the experiment, WA, EOA, EAI and BMD prepared the manuscript.

Acknowledgements and Funding: This study was kindly supported by and conducted at the International Institute of Tropical Agriculture (IITA) Ibadan, Oyo State, Nigeria.

REFERENCES

1. Kotíková Z, Šulc M, Lachman J, Pivec V, Orsák M. and Hamouz K. Carotenoid profile and retention in yellow-, purple- and red-fleshed potatoes after thermal processing. Journal of Food Chemistry, 2016, 197:992–1001.
2. Raposo M, De Morais A and De Morais R. Carotenoids from marine microalgae: A valuable natural source for the prevention of chronic diseases. Journal of Marine Drugs, 2015, 13:5128–5155.
3. Fernández-García E, Carvajal-Lérida I, Jarén-Galán M, Garrido-Fernández J, Pérez-Gálvez A and Hornero-Méndez D. Carotenoids bioavailability from foods: From plant pigments to efficient biological activities. Food Research International, 2012, 46: 438–450.
4. Nishino H, Murakoshi M, Tokuda H and Satomi Y. Cancer prevention by carotenoids. Archives of Biochemistry and Biophysics, 2009, 483:165-168.
5. Kaulmann A and Bohn T. Carotenoids, inflammation, and oxidative stress implications of cellular signaling pathways and relation to chronic disease prevention. Nutrition Research, 2014, 34: 907–929.
6. Schalch W, Cohn W, Barker FM, Kopcke W, Mellerio J, Bird AC, Robson AG, Fitzke FF and van Kuijk FJ. Xanthophyll accumulation in the human retina during supplementation with lutein or zeaxanthin - the LUXEA (LUtein Xanthophyll Eye Accumulation) study. Archives of Biochemistry and Biophysics, 2007, 458(2): 128–135.
7. Britton G and Khachik F. Carotenoid in Food. In: Carotenoids: Nutrition and Health. 5: 3. Britton, G., Liaaen-Jensen, F, Pfander, H (Eds.). Birkhauser Verlag Basel, 2009, ISBN 978-3-7643-7500-3. Pp. 45-66.
8. Rivera S and Canela R. Influence of sample processing on the analysis of carotenoids in maize. Molecules, 2012, 17: 11255-11268.
9. Cazzonelli CI. Carotenoids in nature: Insights from plants and beyond. Functional Plant Biology, 2011, 38: 833–847.
10. Pixley K, Palacios-Rojas N, Babu R, Mutale R, Surles R and Simpungwe E. Biofortification of maize with provitamin A carotenoids. In Tanumihardjo SA, editor. Carotenoids and human health. New York: Springer Science and Business Media. 2013, Pp. 271–92.
11. Scott CE and Eldridge, A. L. Comparison of carotenoid content in fresh, frozen and canned corn. Journal of Food Composition and Analysis, 2005, 18: 551-559.

12. Egesel CO, Wong JC, Lambert RJ, and Rocheford TR. Gene dosage effects on carotenoid concentration in maize grain. Maydica, 2003, 48: 183–190.
13. De Moura FF, Miloff A and Boy E. Retention of provitamin A carotenoids in staple crops targeted for biofortification in Africa: Cassava, maize and sweet potato. Critica Review in Food Science and Nutrition, 2015, 55: 1246–1269.
14. Rodriguez-Amaya DB. Food carotenoids: Chemistry, biology, and technology. UK: John Wiley & Sons, 2015, Pp. 309.
15. Bechoff A, Dhuique-Mayer C, Dornier M, Tomlins KI, Boulanger R, Dufour D, and Westby A. Relationship between the kinetics of β-carotene degradation and formation of norisoprenoids in the storage of dried sweet potato chips. Food Chemistry, 2010, 121: 348-357.
16. Alves RMV, Ito D, Carvalho JLV, Melo W.F, and Godoy RLO. Stability of biofortified sweet potato flour. Brazilian Journal of Food Technology, 2012, 15: 59–71.
17. Ortiz D, Rocheford T and Ferruzi MG. Influence of temperature and humidity on the stability of carotenoids in biofortified maize (Zea mays L.) genotypes during controlled post-harvest storage. Journal of Agriculture and Food Chemistry, 2016, 64: 2727–2736.
18. Gannon B, Kaliwile C, Arscot SA, Schmaelzle S, Chileshe J, Kalungwana N and Tanumiardjo SA. Biofortified orange maize is as efficacious as a vitamin A supplement in Zambian children even in the presence of high liver reserves of vitamin A: A community-based, randomized placebo-controlled trial. American Journal of Clinical Nutrition, 2014, 100: 1541–1550.
19. Mugode L, Ha B, Kaunda A, Sikombe T, Phiri S, Mutale R., Davis C, Tanumihardjo S and De Moura FF. Carotenoid retention of provitamin A maize (Zea mays L.) after Zambian traditional methods of milling, cooking, and storage. Journal of Agriculture and Food Chemistry, 2014, 62: 6317–6325.
20. Awoyale W, Maziya-dixon B, and Menkir A. Retention of pro-vitamin A carotenoids in Ogi powder as affected by packaging materials and storage conditions. Journal of Food and Nutrition Research, 2016, 4(2): 88-93.
21. Awoyale W, Maziya-Dixon B, and Menkir A. Effect of packaging materials and storage conditions on the physicochemical and chemical properties of Ogi powder. Journal of Food, Agriculture and Environment, 2013, 11 (3&4): 242-248.
22. Howe JA and Tanumihardjo SA. Evaluation of analytical methods for carotenoid extraction from biofortified maize (Zea mays sp). Journal of Agriculture and Food Chemistry, 2006, 54: 7992-7997.
23. Taleon V, Mugode L, Cabrera-Soto L and Palacios-Rojas N. Carotenoid retention in biofortified maize using different post-harvest storage and packaging methods. Food Chemistry, 2017, 232: 60–66.
24. Li S, Tayie FAK, Young MF, Rocheford T and White WS. Retention of provitamin A carotenoids in high β-carotene maize (Zea mays) during traditional African household processing. Journal of Agriculture and Food Chemistry, 2007, 55: 10744-10750.
25. Djuikwo VN, Ejoh RA, Gouado I, Mbofung CM and Tanumihardjo SA. Determination of major carotenoids in processed tropical leafy vegetables indigenous to Africa. Food Science and Nutrition, 2011, 2: 793-802.

26. Wu Z, Robinson DS, Hughes RK, Casey R, Hardy D and West SI. Co-oxidation of β-carotene catalyzed by soybean and recombinant pea lipoxygenases. Journal of Agriculture and Food Chemistry, 1999, 47: 4899-4906.
27. Chedea VS and Jisaka M. A review on Lipoxygenase and carotenoids: A co-oxidation story. African Journal of Biotechnology, 2013, 12(20): 2786-2791.
28. Lee DS, Chung SK and Yam KL. The carotenoid loss in dried red pepper products. International Journal of Food Science and Technology, 1992, 27: 179-185.
29. Rao AV and Rao LG. Carotenoids and human health – an Invited review. Pharmacological Research, 2007, 55: 207–216.
30. Zepka LQ and Mercadante AZ. Degradation compounds of carotenoids formed during heating of a simulated cashew apple juice. Food Chemistry, 2009, 117: 28–34.
31. Provesi JG, Dias CO and Amante ER. Changes in carotenoids during processing and storage of pumpkin puree. Food Chemistry, 2011, 128: 195–202.
32. Gama JJT and Sylos CM. Effect of thermal pasteurization and concentration on carotenoid composition of Brazilian valencia orange juice. Food Chemistry, 2007, 100: 1686–1690.
33. Rodriguez-Amaya DB. A guide to carotenoid analysis in foods. Washington, DC: International Life Sciences Institute Press, 1999, Pp. 59.
34. Otten JJ, Hellwig JP and Meyers LD. Dietary Reference Intakes (DRI): The Essential Guide to Nutrient Requirements. Washington, D.C.: The National Academies Press, 2006, Pp. 560.

View publication stats