Int. J. Nonlinear Anal. Appl. In Press, 1–8
ISSN: 2008-6822 (electronic)
http://dx.doi.org/10.22075/ijnaa.2022.26979.3466

Some coefficient problems of a class of close-to-star functions of type α defined by means of a generalized differential operator

Ayotunde Olajide Lasode[a,*], Aminat Olabisi Ajiboye[a], Rasheed Olawale Ayinla[b]

[a]*Department of Mathematics, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria*

[b]*Department of Mathematics and Statistics, Kwara State University, P.M.B. 1530, Malete, Nigeria*

(Communicated by Mugur Alexandru Acu)

Abstract

In this investigation, we studied a class of Bazilevič type close-to-star functions which is defined by a generalized differential operator. The new class generalizes many known and new subclasses of close-to-star functions. Some of the investigated properties are the coefficient bounds and the Fekete-Szegö functional. Our results extend some known and new ones.

Keywords: Analytic function, univalent function, starlike function, close-to-star function of type α, Carathéodory function, Opoola differential operator, coefficient bounds and Fekete-Szegö functional.
2010 MSC: Primary 30C45, Secondary 30C50

1 Introduction and Definitions

In this paper, we let $\mathcal{E} := \{z \in \mathbb{C} : |z| < 1\}$ represent the unit disk and we let $\mathcal{A}$ represent the class of complex-valued functions of the form:

$$f(z) = z + a_2 z^2 + a_3 z^3 + \cdots + a_j z^j + \cdots \quad (z \in \mathcal{E}) \tag{1.1}$$

such that $f(0) = 0 = f'(0) - 1$. $\mathcal{A}$ is called the class of *normalized analytic* functions. Let $\mathcal{S}$, a subset of $\mathcal{A}$, represent the class of functions *analytic* and *univalent* in $\mathcal{E}$.

Also, let $\mathcal{S}^\star$, a subset of $\mathcal{S}$, represent the class of functions of the form:

$$s(z) = z + s_2 z^2 + s_3 z^3 + \cdots + s_j z^j + \cdots \quad (z \in \mathcal{E}) \tag{1.2}$$

such that $\mathcal{R}e(zs'(z)/s(z)) > 0$. The class $\mathcal{S}^\star$ is called the class of *starlike* functions.

Let $\mathcal{C}$ represent the class of analytic functions of the form:

$$c(z) = 1 + c_1 z + c_2 z^2 + c_3 z^3 + \cdots + c_j z^j + \cdots \quad (z \in \mathcal{E}) \tag{1.3}$$

*Corresponding author
Email addresses: lasode_ayo@yahoo.com (Ayotunde Olajide Lasode), amajiboye99@gmail.com (Aminat Olabisi Ajiboye), rasheed.ayinla@kwasu.edu.ng (Rasheed Olawale Ayinla)

Received: April 2022 *Accepted:* June 2022

such that $c(0) = 1$ and $\mathcal{R}e\, c(z) > 0$. The class $\mathcal{C}$ is called the class of *Carathéodory* functions. For additional details on classes $\mathcal{S}$, $\mathcal{S}^\star$ and $\mathcal{C}$, see [9, 21].

A function f in (1.1) is called *close-to-star* if there is a function $s(z) \in \mathcal{S}^\star$ such that

$$\mathcal{R}e\left(\frac{f(z)}{s(z)}\right) > 0 \quad (z \in \mathcal{E}). \tag{1.4}$$

The class of close-to-star was introduced by Reade [15] and it has been studied in various forms by some researchers in [3, 12, 18, 19, 22]. In particular, Babalola et al. [5] studied the class of close-to-star functions of type α defined by using the idea of Bazilevič functions of type α in [20].

Definition 1.1 ([5]). A function $f \in \mathcal{A}$ is called close-to-star of type α if

$$f(z) \in \mathcal{K}^\star(\alpha) := \left\{ f(z) \in \mathcal{A} : \mathcal{R}e\left(\frac{f(z)^\alpha}{s(z)^\alpha}\right) > 0,\ \alpha > 0,\ z \in \mathcal{E} \right\} \tag{1.5}$$

and all powers are regarded as principal determinations only.

Remark 1.2. The following are some remarks on $\mathcal{K}^\star(\alpha)$.

(i) $\mathcal{K}^\star(1)$ is the class introduced by Reade [15].
(ii) $f \in \mathcal{K}^\star(1)$ is not necessarily univalent in $\mathcal{E}$ (see [5, 15]), so also is $f \in \mathcal{K}^\star(\alpha)$.
(iii) The well-known Alexander duality theorem holds. If f is close-to-convex (see [21] for definition), then zf' is close-to-star (see [5, 15]).
(iv) Every starlike function is close-to-star (see [5, 15]).
(v) $\mathcal{R}e(f'(z)/s'(z) > 0 \implies \mathcal{R}e(f(z)/s(z)) > 0$ (see [5, 15, 16]).
(vi) If $f(z) \in \mathcal{K}^*(1)$, then $|a_j| \leqslant j^2$ $(j = \{2, 3, 4, \ldots\})$ (see [15, 16]). Equality occurs for function $f_\delta(z) = (z + \delta z^2)/(1 - \delta z)^3$ $(\delta = \pm i)$ (see [16]).

An interesting aspect of coefficient problems of functions f of form (1.1) is the study of the Fekete-Szegö functional defined by

$$\mathcal{F}(\gamma; f) = |a_3 - \gamma a_2^2|. \tag{1.6}$$

The functional was initiated by Fekete and Szegö [8] and it has received much attention for functions f in class $\mathcal{S}$ and its various subclasses. Some recent investigations in this direction can be found in [1, 4, 7, 11].

In 2017, Opoola [14] introduced a differential operator defined as follows.

Definition 1.3. Let $f \in \mathcal{A}$ be of the form (1.1), then the Opoola differential operator $\mathcal{D}^{n,\beta,\eta,\tau} : \mathcal{A} \longrightarrow \mathcal{A}$ is defined by

$$\begin{aligned}
\mathcal{D}^{0,\beta,\eta,\tau} f(z) &= f(z) \\
\mathcal{D}^{1,\beta,\eta,\tau} f(z) &= (1 + (\beta - \eta - 1)\tau) f(z) - z\tau(\beta - \eta) + z\tau f'(z) = \Delta_\tau f(z) \\
\mathcal{D}^{2,\beta,\eta,\tau} f(z) &= \Delta_\tau(\mathcal{D}^{1,\beta,\eta,\tau} f(z)) \\
\mathcal{D}^{3,\beta,\eta,\tau} f(z) &= \Delta_\tau(\mathcal{D}^{2,\beta,\eta,\tau} f(z)) \\
&\ \ \vdots \qquad \vdots \qquad \vdots \\
\mathcal{D}^{n,\beta,\eta,\tau} f(z) &= \Delta_\tau(\mathcal{D}^{n-1,\beta,\eta,\tau} f(z))
\end{aligned}$$

which implies that

$$\mathcal{D}^{n,\beta,\eta,\tau} f(z) = z + \sum_{j=2}^{\infty} (1 + (j + \beta - \eta - 1)\tau)^n a_j z^j \tag{1.7}$$

or for brevity,

$$\mathcal{D}^{n,\beta,\eta,\tau} f(z) = z + \sum_{j=2}^{\infty} \Lambda_j a_j z^j \quad (z \in \mathcal{E}) \tag{1.8}$$

where $\Lambda_j = (1 + (j + \beta - \eta - 1)\tau)^n$, $\tau \geqslant 0$, $0 \leqslant \eta \leqslant \beta$ and $n \in \mathbb{N} \cup \{0\}$.

Remark 1.4. The following properties hold in (1.7).

(i) $\mathcal{D}^{0,\beta,\eta,\tau}f(z) = \mathcal{D}^{n,\beta,\eta,0}f(z) = \mathcal{D}^{0,\beta,\eta,0}f(z) = f(z) \in \mathcal{A}$ in (1.1).
(ii) $\mathcal{D}^{n,\beta,\beta,1}f(z) = \mathcal{D}^{n,\eta,\eta,1}f(z) = \mathcal{D}^{n}f(z)$ is the Sălăgean differential operator introduced in [17].
(iii) $\mathcal{D}^{n,\beta,\beta,\tau}f(z) = \mathcal{D}^{n,\eta,\eta,\tau}f(z) = \mathcal{D}^{n,\tau}f(z)$ is the Al-Oboudi differential operator introduced in [2].

The new class investigated in this paper is therefore defined as follows.

Definition 1.5. If $n \in \mathbb{N} \cup \{0\}$, $0 \leqslant \eta \leqslant \beta$ and $\tau \geqslant 0$, then a function $f(z) \in \mathcal{A}$ is said to be a member of class $\mathcal{K}^{\star}(n,\beta,\eta,\tau;\alpha)$ if it satisfies the geometric condition

$$\mathcal{R}e\left(\frac{(\mathcal{D}^{n,\beta,\eta,\tau}f(z))^{\alpha}}{(\mathcal{D}^{n,\beta,\eta,\tau}s(z))^{\alpha}}\right) > 0 \quad (\alpha \geqslant 1,\ z \in \mathcal{E}) \tag{1.9}$$

where $\mathcal{D}^{n,\beta,\eta,\tau}$ is the Opoola differential operator in (1.7) and all powers are regarded as principal determinations only.

Remark 1.6. Note that $\mathcal{K}^{\star}(n,\beta,\eta,0;\alpha) = \mathcal{K}^{\star}(0,\beta,\eta,\tau;\alpha) = \mathcal{K}^{\star}(0,\beta,\eta,0;\alpha) = \mathcal{K}^{\star}(\alpha)$ is the class studied by Babalola et al. [5]. And that $\mathcal{K}^{\star}(n,\beta,\eta,0;1) = \mathcal{K}^{\star}(0,\beta,\eta,\tau;1) = \mathcal{K}^{\star}(0,\beta,\eta,0;1) = \mathcal{K}^{\star}(1)$ is the class studied by Reade [15].

It is interesting to know that class $\mathcal{K}^{\star}(n,\beta,\eta,\tau;\alpha)$ is non-empty as shown in the following examples.

Example 1.7. *If $s(z) = z$, then for $f(z) \in \mathcal{A}$ of the form (1.1),*

$$\begin{aligned}
\mathcal{D}^{n,\beta,\eta,\tau}f_1(z) &= \{z^{\alpha}(1+z)\}^{\frac{1}{\alpha}} = z + \lambda(\alpha,1)z^2 + \lambda(\alpha,2)z^3 + \lambda(\alpha,3)z^4 + \lambda(\alpha,4)z^5 + \cdots,\\
\mathcal{D}^{n,\beta,\eta,\tau}f_2(z) &= \{z^{\alpha}(1-z)\}^{\frac{1}{\alpha}} = z - \lambda(\alpha,1)z^2 + \lambda(\alpha,2)z^3 - \lambda(\alpha,3)z^4 + \lambda(\alpha,4)z^5 - \cdots,\\
\mathcal{D}^{n,\beta,\eta,\tau}f_3(z) &= \{z^{\alpha}(1+z)^{-1}\}^{\frac{1}{\alpha}} = z + \lambda(\alpha,1)z^2 + \lambda(\alpha,2)z^3 + \lambda(\alpha,3)z^4 + \lambda(\alpha,4)z^5 + \cdots,\\
\mathcal{D}^{n,\beta,\eta,\tau}f_4(z) &= \{z^{\alpha}(1-z)^{-1}\}^{\frac{1}{\alpha}} = z - \lambda(\alpha,1)z^2 + \lambda(\alpha,2)z^3 - \lambda(\alpha,3)z^4 + \lambda(\alpha,4)z^5 - \cdots,
\end{aligned}$$

and

$$\begin{aligned}
\mathcal{D}^{n,\beta,\eta,\tau}f_5(z) = \left\{z^{\alpha}\frac{1+z}{1-z}\right\}^{\frac{1}{\alpha}} &= z + 2\lambda(\alpha,1)z^2 + [2\lambda(\alpha,1) + 4\lambda(\alpha,2)]z^3 + [2\lambda(\alpha,1) + 8\lambda(\alpha,2) + 8\lambda(\alpha,3)]z^4\\
&\quad + [2\lambda(\alpha,1) + 12\lambda(\alpha,2) + 24\lambda(\alpha,3) + 16\lambda(\alpha,4)]z^5 + \cdots
\end{aligned}$$

are in class $\mathcal{K}^{\star}(n,\beta,\eta,\tau;\alpha)$ such that for $m \in \mathbb{N}$,

$$\lambda(\alpha,m) = \begin{cases} \frac{1}{m!}\prod_{t=0}^{m-1}\left(\frac{1}{\alpha} - t\right) & for\ power \quad \frac{1}{\alpha} > 0\\ \frac{(-1)^m}{m!}\prod_{t=0}^{m-1}\left(\frac{1}{\alpha} + t\right) & for\ power \quad \frac{1}{\alpha} < 0. \end{cases}$$

Proof . From (1.9), we can say that

$$\frac{(\mathcal{D}^{n,\beta,\eta,\tau}f_k(z))^{\alpha}}{(\mathcal{D}^{n,\beta,\eta,\tau}s(z))^{\alpha}} = \begin{cases} 1+z & for \quad k=1\\ 1-z & for \quad k=2\\ \frac{1}{1+z} & for \quad k=3\\ \frac{1}{1-z} & for \quad k=4\\ \frac{1+z}{1-z} & for \quad k=5 \end{cases} \tag{1.10}$$

where all functions on the RHS of (1.10) are well-known functions in class $\mathcal{C}$ defined above. Now $s(z) = z \implies (\mathcal{D}^{n,\beta,\eta,\tau}s(z))^{\alpha} = z^{\alpha}$ and $(\mathcal{D}^{n,\beta,\eta,\tau}f_k(z))^{\alpha}$ $(k = \{1,2,3,4,5\})$ in (1.10) give the functions in Examples 1.7 by simple calculation. This completes the proof. □

2 Relevant Lemmas

Lemma 2.1 ([21]). If $c(z) \in \mathcal{C}$, then $|c_j| \leqslant 2 \quad (j \in \mathbb{N})$.

Lemma 2.2 ([6]). If $c(z) \in \mathcal{C}$ and $u \in \mathbb{R}$, then

$$\left|c_2 - u\frac{c_1^2}{2}\right| \leqslant \begin{cases} 2(1-u) & for \quad u \leqslant 0, \\ 2 & for \quad 0 \leqslant u \leqslant 2, \\ 2(u-1) & for \quad u \geqslant 2. \end{cases}$$

Lemma 2.3 ([13]). If $c(z) \in \mathcal{C}$ and $i, j \in \mathbb{N}$, then $|c_{i+j} - uc_ic_j| \leqslant 2$ for $0 \leqslant u \leqslant 1$.

Lemma 2.4 ([21]). If $s(z) \in \mathcal{S}^\star$, then $|s_j| \leqslant j \quad (j \in \mathbb{N} \setminus \{1\})$.

Lemma 2.5 ([10]). If $s(z) \in \mathcal{S}^\star$ and $\rho \in \mathbb{R}$, then

$$|s_3 - \rho s_2^2| \leqslant \begin{cases} 3 - 4\rho & for \quad \rho \leqslant \frac{1}{2} \\ 1 & for \quad \frac{1}{2} \leqslant \rho \leqslant 1 \\ 4\rho - 3 & for \quad \rho \geqslant 1. \end{cases}$$

3 Main Results

In what follows, let $n \in \mathbb{N} \cup \{0\}$, $0 \leqslant \eta \leqslant \beta$, $\tau \geqslant 0$ and $\alpha \geqslant 1$ throughout this work unless otherwise mentioned. The following theorems are the results obtained.

Theorem 3.1. If $f(z) \in \mathcal{K}^\star(n, \beta, \eta, \tau; \alpha)$, then

$$|a_2| \leqslant \frac{2}{\alpha\Lambda_2}\left\{1 + \alpha\Lambda_2\right\}, \tag{3.1}$$

$$|a_3| \leqslant \frac{2}{\alpha\Lambda_3}\left\{1 + 2\Lambda_2 + \frac{3\alpha\Lambda_3}{2}\right\}, \tag{3.2}$$

$$|a_4| \leqslant \frac{2}{\alpha\Lambda_4}\left\{1 + 2\Lambda_2 + 3\Lambda_3 + 2\alpha\Lambda_4 + \frac{2(\alpha-1)(2\alpha-1)}{3\alpha^2}\right\}, \tag{3.3}$$

$$\begin{aligned} |a_5| \leqslant \frac{2}{\alpha\Lambda_5}\Bigg\{1 + 2\Lambda_2 + 3\Lambda_3 + 2(\alpha-1)(2\alpha-1) + 2[2\Lambda_4 + 3(\alpha-1)|4-3\alpha|\Lambda_2\Lambda_3] \\ + \frac{4(\alpha-1)|\alpha-2||3\alpha-4|\Lambda_2}{3} + \frac{(\alpha-1)}{\alpha} + \frac{\alpha|\alpha-2|}{4}[16\Lambda_2\Lambda_4 + 9\Lambda_3^2] + \frac{5\alpha\Lambda_5}{2}\Bigg\}. \end{aligned} \tag{3.4}$$

Proof . The geometric expression in (1.9) can be expressed as

$$\frac{(\mathcal{D}^{n,\beta,\eta,\tau}f(z))^\alpha}{z^\alpha} = \frac{(\mathcal{D}^{n,\beta,\eta,\tau}s(z))^\alpha}{z^\alpha}c(z). \tag{3.5}$$

Using (1.1), (1.8) and binomially expanding LHS of (3.5) we obtain

$$\begin{aligned} \frac{(\mathcal{D}^{n,\beta,\eta,\tau}f(z))^\alpha}{z^\alpha} &= 1 + \alpha\Lambda_2a_2z \\ &\quad + \left\{\alpha\Lambda_3a_3 + \frac{\alpha(\alpha-1)}{2!}\Lambda_2^2a_2^2\right\}z^2 \\ &\quad + \left\{\alpha\Lambda_4a_4 + \frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3a_2a_3 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3a_2^3\right\}z^3 \\ &\quad + \Bigg\{\alpha\Lambda_5a_5 + \frac{\alpha(\alpha-1)}{2!}(2\Lambda_2\Lambda_4a_2a_4 + \Lambda_3^2a_3^2) + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}3\Lambda_2^2\Lambda_3a_2^2a_3 \\ &\qquad + \frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!}\Lambda_2^4a_2^4\Bigg\}z^4 \\ &\quad + \cdots . \end{aligned} \tag{3.6}$$

And using (1.2), (1.3), (1.8) and binomially expanding RHS of (3.5) we obtain

$$\begin{aligned}\frac{(\mathcal{D}^{n,\beta,\eta,\tau}s(z))^{\alpha}}{z^{\alpha}}c(z) &= 1+\{c_1+\alpha\Lambda_2 s_2\}z\\ &+\left\{c_2+\alpha\Lambda_2 s_2c_1+\alpha\Lambda_3 s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2\right\}z^2\\ &+\left\{c_3+\alpha\Lambda_2 s_2c_2+\left(\alpha\Lambda_3 s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2\right)c_1\right.\\ &\quad\left.+\frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3 s_2s_3+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3s_2^3+\alpha\Lambda_4 s_4\right\}z^3\\ &+\left\{c_4+\alpha\Lambda_2 s_2c_3+\left(\alpha\Lambda_3 s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2\right)c_2\right.\\ &\quad+\left(\alpha\Lambda_4 s_4+\frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3 s_2s_3+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3s_2^3\right)c_1\\ &\quad+\frac{\alpha(\alpha-1)}{2!}(2\Lambda_2\Lambda_4 s_2s_4+\Lambda_3^2s_3^2)+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}(3\Lambda_2^2\Lambda_3 s_2^2s_3)\\ &\quad\left.+\frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!}\Lambda_2^4s_2^4+\alpha\Lambda_5 s_5\right\}z^4\\ &+\cdots. \end{aligned}\tag{3.7}$$

Now, if we equate the coefficients in (3.6) and (3.7), then we obtain

$$\alpha\Lambda_2a_2=c_1+\alpha\Lambda_2s_2,\tag{3.8}$$

$$\alpha\Lambda_3a_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2a_2^2=c_2+\alpha\Lambda_2s_2c_1+\alpha\Lambda_3s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2,\tag{3.9}$$

$$\begin{aligned}\alpha\Lambda_4a_4+\frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3a_2a_3+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3a_2^3 &= c_3+\alpha\Lambda_2s_2c_2\\ &+\left(\alpha\Lambda_3s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2\right)c_1+\frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3s_2s_3+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3s_2^3+\alpha\Lambda_4s_4\end{aligned}\tag{3.10}$$

and

$$\begin{aligned}\alpha\Lambda_5a_5+\frac{\alpha(\alpha-1)}{2!}(2\Lambda_2\Lambda_4a_2a_4+\Lambda_3^2a_3^2)+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}3\Lambda_2^2\Lambda_3a_2^2a_3+\frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!}\Lambda_2^4a_2^4\\ =c_4+\alpha\Lambda_2s_2c_3+\left(\alpha\Lambda_3s_3+\frac{\alpha(\alpha-1)}{2!}\Lambda_2^2s_2^2\right)c_2\\ +\left(\alpha\Lambda_4s_4+\frac{\alpha(\alpha-1)}{2!}2\Lambda_2\Lambda_3s_2s_3+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}\Lambda_2^3s_2^3\right)c_1\\ +\frac{\alpha(\alpha-1)}{2!}(2\Lambda_2\Lambda_4s_2s_4+\Lambda_3^2s_3^2)+\frac{\alpha(\alpha-1)(\alpha-2)}{3!}3\Lambda_2^2\Lambda_3s_2^2s_3+\frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!}\Lambda_2^4s_2^4+\alpha\Lambda_5s_5.\end{aligned}\tag{3.11}$$

Simple calculation shows that (3.8) leads to

$$a_2=\frac{1}{\alpha\Lambda_2}c_1+s_2\tag{3.12}$$

so that by applying triangle inequality and Lemmas 2.1 and 2.4 in (3.12) lead to inequality (3.1).

Using (3.12) in (3.9) leads to

$$a_3=\frac{1}{\alpha\Lambda_3}c_2+\frac{\Lambda_2s_2}{\alpha\Lambda_3}c_1-\frac{\alpha-1}{2\alpha^2\Lambda_3}c_1^2+s_3\tag{3.13}$$

so that by applying triangle inequality we obtain

$$|a_3|=\left|\frac{1}{\alpha\Lambda_3}c_2+\frac{\Lambda_2s_2}{\alpha\Lambda_3}c_1-\frac{\alpha-1}{2\alpha^2\Lambda_3}c_1^2+s_3\right|\leqslant\frac{1}{\alpha\Lambda_3}\left\{\left|c_2-\frac{\alpha-1}{\alpha}\frac{c_1^2}{2}\right|+\Lambda_2|s_2||c_1|+\alpha\Lambda_3|s_3|\right\}$$

and using Lemmas 2.1, 2.2 and 2.4 we obtain (3.2).

Using (3.12) and (3.13) in (3.10) leads to

$$a_4 = \frac{1}{\alpha\Lambda_4}c_3 + \frac{\Lambda_2 s_2}{\alpha\Lambda_4}c_2 + \frac{\Lambda_3 s_3}{\alpha\Lambda_4}c_1 - \frac{\alpha-1}{\alpha^2\Lambda_4}c_1c_2 - \frac{(\alpha-1)\Lambda_2 s_2}{2\alpha^2\Lambda_4}c_1^2 + \frac{(\alpha-1)(2\alpha-1)}{6\alpha^3\Lambda_4}c_1^3 + s_4 \tag{3.14}$$

so that by applying triangle inequality,

$$\begin{aligned}|a_4| &= \left|\frac{1}{\alpha\Lambda_4}c_3 + \frac{\Lambda_2 s_2}{\alpha\Lambda_4}c_2 + \frac{\Lambda_3 s_3}{\alpha\Lambda_4}c_1 - \frac{\alpha-1}{\alpha^2\Lambda_4}c_1c_2 - \frac{(\alpha-1)\Lambda_2 s_2}{2\alpha^2\Lambda_4}c_1^2 + \frac{(\alpha-1)(2\alpha-1)}{6\alpha^3\Lambda_4}c_1^3 + s_4\right| \\ &\leqslant \frac{1}{\alpha\Lambda_4}\left\{\left|c_3 - \frac{\alpha-1}{\alpha}c_1c_2\right| + \Lambda_2|s_2|\left|c_2 - \frac{\alpha-1}{\alpha}\frac{c_1^2}{2}\right| + \Lambda_3|s_3||c_1| + \frac{(\alpha-1)(2\alpha-1)}{6\alpha^2}|c_1|^3 + \alpha\Lambda_4|s_4|\right\}\end{aligned}$$

and using Lemmas 2.1, 2.2, 2.3 and 2.4 we obtain (3.3).

Using (3.12), (3.13) and (3.14) in (3.11) leads to

$$\begin{aligned}a_5 = \frac{1}{\alpha\Lambda_5}\Bigg\{&c_4 + \Lambda_2 s_2 c_3 + \Lambda_3 s_3 c_2 + [\Lambda_4 s_4 + (\alpha-1)(4-3\alpha)\Lambda_2\Lambda_3 s_2 s_3]c_1 \\ &- \frac{(\alpha-1)\Lambda_3 s_3}{2\alpha}c_1^2 + \frac{(\alpha-1)(\alpha-2)(3\alpha-4)\Lambda_2 s_2}{6\alpha^2}c_1^3 - \frac{(\alpha-1)(2\alpha-1)}{24\alpha^3}c_1^4 \\ &- \frac{(\alpha-1)\Lambda_2 s_2}{\alpha}c_1c_2 - \frac{(\alpha-1)}{\alpha}c_1c_3 + \frac{(\alpha-1)(2\alpha-1)}{2\alpha^2}c_1^2c_2 - \frac{(\alpha-1)}{2\alpha}c_2^2 \\ &+ \frac{\alpha(\alpha-2)}{2}[2\Lambda_2\Lambda_4 s_2 s_4 + \Lambda_3^2 s_3^2] + \alpha\Lambda_5 s_5\Bigg\}\end{aligned}$$

and

$$\begin{aligned}|a_5| = \frac{1}{\alpha\Lambda_5}\Bigg|&\left(c_4 - \frac{(\alpha-1)}{\alpha}c_1c_3\right) + \left(\Lambda_2 s_2 c_3 - \frac{(\alpha-1)\Lambda_2 s_2}{\alpha}c_1c_2\right) + \left(\Lambda_3 s_3 c_2 - \frac{(\alpha-1)\Lambda_3 s_3}{2\alpha}c_1^2\right) \\ &+ \left(\frac{(\alpha-1)(2\alpha-1)}{2\alpha^2}c_1^2c_2 - \frac{(\alpha-1)(2\alpha-1)^2}{24\alpha^3}c_1^4\right) + \frac{(\alpha-1)(\alpha-2)(3\alpha-4)\Lambda_2 s_2}{6\alpha^2}c_1^3 \\ &+ [\Lambda_4 s_4 + (\alpha-1)(4-3\alpha)\Lambda_2\Lambda_3 s_2 s_3]c_1 - \frac{(\alpha-1)}{2\alpha}c_2^2 \\ &+ \frac{\alpha(\alpha-2)}{2}[2\Lambda_2\Lambda_4 s_2 s_4 + \Lambda_3^2 s_3^2] + \alpha\Lambda_5 s_5\Bigg|\end{aligned}$$

so that by applying triangle inequality we obtain

$$\begin{aligned}|a_5| \leqslant \frac{1}{\alpha\Lambda_5}\Bigg\{&\left|c_4 - \frac{(\alpha-1)}{\alpha}c_1c_3\right| + \Lambda_2|s_2|\left|c_3 - \frac{(\alpha-1)}{\alpha}c_1c_2\right| + \Lambda_3|s_3|\left|c_2 - \frac{(\alpha-1)}{2\alpha}c_1^2\right| \\ &+ \frac{(\alpha-1)(2\alpha-1)}{2\alpha^2}|c_1|^2\left|c_2 - \frac{(2\alpha-1)}{6\alpha}\frac{c_1^2}{2}\right| + \frac{(\alpha-1)|\alpha-2||3\alpha-4|\Lambda_2|s_2|}{6\alpha^2}|c_1|^3 \\ &+ [\Lambda_4|s_4| + (\alpha-1)|4-3\alpha|\Lambda_2\Lambda_3|s_2||s_3|]|c_1| + \frac{(\alpha-1)}{2\alpha}|c_2|^2 \\ &+ \frac{\alpha(\alpha-2)}{2}[2\Lambda_2\Lambda_4|s_2||s_4| + \Lambda_3^2|s_3|^2] + \alpha\Lambda_5|s_5|\Bigg\}\end{aligned}$$

and using Lemmas 2.1, 2.2, 2.3 and 2.4 we obtain (3.4). □

Remark 3.2. Setting $n = 0$ (or $\tau = 0$) makes inequalities (3.1), (3.2) and (3.3) to become the results of Babalola et al [5].

Theorem 3.3. If $f(z) \in \mathcal{K}^{\star}(n,\beta,\eta,\tau;\alpha)$, then for $x \in \mathbb{R}$,

$$|a_3 - xa_2^2| \leqslant \begin{cases} \frac{4}{\alpha\Lambda_3}\left(1 - \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\right) & for \quad x \leqslant \frac{(1-\alpha)\Lambda_2^2}{2\Lambda_3}, \\ \frac{4}{\alpha\Lambda_3} & for \quad \frac{(1-\alpha)\Lambda_2^2}{2\Lambda_3} \leqslant x \leqslant \frac{2\alpha\Lambda_2^2+(1-\alpha)\Lambda_2^2}{2\Lambda_3}, \\ \frac{4}{\alpha\Lambda_3}\left(\frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2} - 1\right) & for \quad x \geqslant \frac{2\alpha\Lambda_2^2+(1-\alpha)\Lambda_2^2}{2\Lambda_3}, \\ \frac{2}{\alpha\Lambda_3}\left(\frac{4|\Lambda_2^2-2\Lambda_3 x|}{\Lambda_2} + \alpha\Lambda_3(3-4x)\right) & for \quad x \leqslant \frac{1}{2}, \\ \frac{2}{\alpha\Lambda_3}\left(\frac{4|\Lambda_2^2-2\Lambda_3 x|}{\Lambda_2} + \alpha\Lambda_3\right) & for \quad \frac{1}{2} \leqslant x \leqslant 1, \\ \frac{2}{\alpha\Lambda_3}\left(\frac{4|\Lambda_2^2-2\Lambda_3 x|}{\Lambda_2} + \alpha\Lambda_3(4x-3)\right) & for \quad x \geqslant 1. \end{cases} \tag{3.15}$$

Proof . Consider (3.12) and (3.13) in (1.6) and for $x \in \mathbb{R}$ implies that

$$\begin{aligned} |a_3 - xa_2^2| &= \left|\frac{1}{\alpha\Lambda_3}c_2 + \frac{\Lambda_2 s_2}{\alpha\Lambda_3}c_1 - \frac{\alpha-1}{2\alpha^2\Lambda_3}c_1^2 + s_3 - x\left(\frac{1}{\alpha\Lambda_2}c_1 + s_2\right)^2\right| \\ &= \frac{1}{\alpha\Lambda_3}\left|c_2 - \left(\frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\right)\frac{c_1^2}{2} + \frac{(\Lambda_2^2-2\Lambda_3 x)s_2}{\Lambda_2}c_1 + \alpha\Lambda_3(s_3 - xs_2^2)\right| \\ &\leqslant \frac{1}{\alpha\Lambda_3}\left|c_2 - \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\frac{c_1^2}{2}\right| + \frac{1}{\alpha\Lambda_3}\left|\frac{\Lambda_2^2-2\Lambda_3 x s_2}{\Lambda_2}c_1 + \alpha\Lambda_3(s_3 - xs_2^2)\right|. \end{aligned}$$

It is easy to see that

$$|a_3 - xa_2^2| \leqslant \frac{2}{\alpha\Lambda_3}\left|c_2 - \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\frac{c_1^2}{2}\right| \quad \text{if} \quad \left|c_2 - \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\frac{c_1^2}{2}\right| \geqslant \left|\frac{\Lambda_2^2-2\Lambda_3 x s_2}{\Lambda_2}c_1 + \alpha\Lambda_3(s_3 - xs_2^2)\right| \tag{3.16}$$

and

$$|a_3 - xa_2^2| \leqslant \frac{2}{\alpha\Lambda_3}\left|\frac{\Lambda_2^2-2\Lambda_3 x s_2}{\Lambda_2}c_1 + \alpha\Lambda_3(s_3 - xs_2^2)\right| \quad \text{if} \quad \left|c_2 - \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}\frac{c_1^2}{2}\right| \leqslant \left|\frac{\Lambda_2^2-2\Lambda_3 x s_2}{\Lambda_2}c_1 + \alpha\Lambda_3(s_3 - xs_2^2)\right|. \tag{3.17}$$

Applying Lemma 2.2 in (3.16) shows that

$$|a_3 - xa_2^2| \leqslant \frac{2}{\alpha\Lambda_3}\left|c_2 - u\frac{c_1^2}{2}\right| \quad \text{where } u = \frac{(\alpha-1)\Lambda_2^2+2\Lambda_3 x}{\alpha\Lambda_2^2}$$

so that for u in the intervals $u \leqslant 0$, $0 \leqslant u \leqslant 2$ and $u \geqslant 2$ we obtain the first three results in (3.15).

On the other hand, (3.17) simplifies to

$$|a_3 - xa_2^2| \leqslant \frac{2}{\alpha\Lambda_3}\left\{\frac{|\Lambda_2^2-2\Lambda_3 x||s_2|}{\Lambda_2}|c_1| + \alpha\Lambda_3|s_3 - xs_2^2|\right\}$$

so that by applying Lemmas 2.1, 2.4 and 2.5 we obtain the last three results in (3.15) and the proof is complete. □

Acknowledgments

The authors sincerely appreciate the referees' time taken to review and give constructive comments which improved this research.

References

[1] A.O. Ajiboye and K.O. Babalola, *Linear sum of analytic functions defined by a convolution operator,* J. Nigerian Math. Soc. **39** (2020), 173–182.

[2] F.M. Al-Oboudi, *On univalent functions defined by a generalised Sălăgean operator*, Intern. J. Math. Math. Sci. **2004** (2004), 1429–1436.

[3] M. Arif, J. Dziok, M. Raza and J. Sokól, *On products of multivalent close-to-star functions*, J. Inequal. Appl. **5** (2015), 1–14.

[4] R.O. Ayinla and T.O. Opoola, *The Fekete Szegö functional and second Hankel determinant for a certain subclass of analytic functions*, Appl. Math. **10** (2019), 1071–1078.

[5] K.O. Babalola, A.O. Olasupo and C.N. Ejieji, *Early coefficients of close-to-star functions of type* α, J. Nigerian Math. Soc. **31** (2012), 185–189.

[6] K.O. Babalola and T.O. Opoola, *On the coefficients of a certain class of analytic functions,* Adv. Inequal. Ser. **1** (2008), 1–13.

[7] R.A. Bello and T.O. Opoola, *Upper bounds for Fekete-Szegö functions and the second Hankel determinants for a class of starlike functions,* Intern. J. Math. **13** (2017), 34–39.

[8] M. Fekete and G. Szegö, *Eine bemerkung über ungerade schlichte funktionen*, J. Lond. Math. Soc. **8** (1933), 85–89.

[9] A.A. James, A.O. Lasode and B.O. Moses, *Geometric conditions for starlikeness and convexity of univalent functions*, IOSR J. Math. **3** (2012), 15–23.

[10] F.R. Keogh and E.P. Merkes, *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12.

[11] A.O. Lasode and T.O. Opoola, *Fekete-Szegö estimates and second Hankel determinant for a generalized subfamily of analytic functions defined by q-differential operator*, Gulf J. Math. **11** (2021), 36–43.

[12] A. Lecko and Y.J. Sim, *Coefficient problems in the subclasses of close-to-star functions,* Results Math. **74** (2019), 1–14.

[13] A.E. Livingston, *The coefficients of multivalent close-to-convex functions*, Proc. Amer. Math. Soc. **21** (1969), 545–552.

[14] T.O. Opoola, *On a subclass of univalent functions defined by a generalised differential operator*, Intern. J. Math. Anal. **11** (2017), 869–876.

[15] M. Reade, *On close-to-convex univalent functions*, Mich. Math. J. **3** (1955), 59-–62.

[16] M.S. Robertson, *Analytic functions star-like in one direction,* Amer. J. Math. **58** (1936), 465–472.

[17] G.S. Sălăgean, *Subclasses of univalent functions*, Lect. Notes Math. **1013** (1983), 362–372.

[18] C. Selvaraj and S. Logu, *A subclass of close-to-star functions,* Malay. J. Math. **5** (2019), 290–295.

[19] A. Sen, M. Aydoğan and Y. Polatoğlu, *Distortion estimate and the radius of starlikeness of Janowski close-to-star functions,* Theory Appl. Math. Comput. Sci. **1** (2011), 89–92.

[20] R. Singh, *On Bazilevič functions,* Proc. Amer. Math. Soc. **38** (1973), 261–271.

[21] D. K. Thomas, N. Tuneski and A. Vasudevarao, *Univalent Functions: A Primer,* Walter de Gruyter Inc., Berlin, 2018.

[22] P. Zaprawa, *On close-to-star functions,* Bull. Belg. Math. Soc. **16** (2009), 469–480.