IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Common Coupled Fixed Point Theorems without Compatibility in Partially Ordered Metric Spaces

K. R. Tijani[1*], O. T. Wahab[2], I. F. Usamot[3], S. M. Alata[4]

1 Department of Mathematical Sciences, Osun State University, Osogbo, Nigeria
2 Department of Mathematics, Kwara State University, Malete, Nigeria
3 Department of Mathematics, University of Ilorin, Ilorin, Nigeria
4 Department of Computer Science, KwaraCAILS, Ilorin, Nigeria
* Corresponding author: kamiludeen.tijani@uniosun.edu.ng, kkrotimi72@gmail.com,kamil_tijani2000@yahoo.com

Article Info
Received: 22 February 2022 Revised: 15 June 2022
Accepted: 15 June 2022 Available online: 30 June 2022

Abstract

A perfect blend of requirements for the proof of common coupled fixed point theorems in partially ordered metric space without the assumptions of (weak) compatibility is accomplished. Previous attempts in this direction involving these assumptions mostly ensure existence of coupled coincidence points. In many existing works in this area, attempt have been made to prove the existence of common coupled fixed points. However, only identity mappings can satisfy the conditions of the theorems. The method of proof presented in this present work is powerful in view of the fact that it guarantees the existence of common coupled fixed points without the imposition of (weak) compatibility conditions and identity mappings. To illustrate the results, an example is provided.

Keywords: Coupled fixed point of mappings, Partially ordered metric spaces, Common coupled fixed points, Identity mappings, Compatibility, Mixed monotone properties.
MSC2010: 37C25

1 Introduction

The following definitions are to be employed in the sequel.
Definition 1.1 [1–3] Let (X, d) be a metric space and let $T : X \times X \to X$ be a mapping. An element $(a, b) \in X \times X$ is said to be a *coupled fixed point* of T if $a = T(a, b)$ and $b = T(b, a)$.
Definition 1.2. [4, 5] Let (X, d) be a metric space and let $T : X \times X \to X$, $S : X \times X \to X$ be mappings. An element $(a, b) \in X \times X$ is said to be a *common coupled fixed point* of T and S if $a = S(a, b) = T(a, b)$ and $b = S(b, a) = T(b, a)$.
Definition 1.3 [9–11]: Consider a function $\psi : {}^{+} \to {}^{+}$ satisfying
(i) ψ is monotone increasing;
(ii) $\psi^n(t) \to 0$, as n $\to \infty$;

This work is licensed under a Creative Commons Attribution 4.0 International License.

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

(iii) $\sum_{n=0}^{\infty} \psi^n(t)$ converges for all $t > 0$.

(1) A function ψ satisfying (i) and (ii) above is called a *comparison function.*
(2) A function ψ satisfying (i) and (iii) above is called a *(c)-comparison function.*

Remark 1.4 [10, 11] :
(i) Any (c)-comparison function is a comparison function.
(ii) Every comparison function satisfies $\psi(0) = 0$.

The notion of coupled fixed points was introduced by Chang and Ma [13]. Since then, the concept has been of great interest to many researchers in the metrical fixed point theory.
Bhaskar and Lakshmikantham [1] proved a coupled fixed point theorem in a metric space endowed with partial order.
They introduced the following notions of a mapping satisfying the mixed monotone property:
Definition 1.5 [1]: Let $(X, \leq)$ be a partially ordered set and $F : X \times X \to X$, a mapping. The mapping F is said to have the mixed monotone property if F is monotone non-decreasing in its first argument and is monotone non-increasing in its second argument, that is, for any a, $b \in X$,

$$a_1, a_2 \in X, \quad a_1 \leq a_2 \Rightarrow F(a_1, b) \leq F(a_2, b)$$

and

$$b_1, b_2 \in X, \quad b_1 \leq b_2 \Rightarrow F(a, b_1) \geq F(a, b_2).$$

They proved the following theorem, considering a function satisfying the mixed monotone property:
Theorem 1.6: [1] Let $F : X \times X \to X$ be a continuous mapping having the mixed monotone property on X. Assume that there exists a $k \in [0, 1)$ with

$$d(F(a, b), F(u, v)) \leq \frac{k}{2}[d(a, u) + d(b, v)], \forall\ a \geq u, b \leq v.$$

If there exists $a_0, b_0 \in X$ such that

$$a_0 \leq F(a_0, b_0) \quad \textit{and} \quad b_0 \geq F(b_0, a_0),$$

then, there exist a, $b \in X$ such that $a = F(a, b) \quad \textit{and} \quad b = F(b, a)$.
The result has been generalised and extended by Lakshmikantam and Ciric [18], by introducing mixed g-monotone mapping and proving coupled coincidence and coupled common fixed point theorems for such contractive mappings in partially ordered metric spaces.

Several other authors have generalized the results of Bhaskar and Lakshmikantham [1]. Also, for more results on coupled fixed point theorems, interested readers can check [2,3,8,12,14,20–24,26–28] and others in the literature.

The problem of proving common coupled fixed point theorems in the setting of partially ordered metric spaces using the mixed monotone property has been of immense research interest. Lakshmikantam and Ciric [18] established common coupled fixed point theorems by using g-mixed monotone property while one of the mappings is made a proper subset of the other. They proved the existence of coupled coincidence points while coupled fixed point can be established only by assumption of identity on one of the mappings.
In the same line of argument, Abbas *et al.* [4] and Kim and Chandok [17] proved coupled coincidence points of mappings using the mixed monotone property and making one of the mappings a proper subset of the other.
Moerover, Abbas *et al.* [5] proved coupled coincidence point and common coupled fixed points for w-compatible mappings in partially ordered metric spaces. Also Kadelburg *et al.* [16] proved common coupled fixed points for compatible mappings in partially ordered metric spaces. Several researchers have proved common coupled fixed point theorems employing the notions mentioned above. For details on the common coupled fixed point theorems in partially ordered metric spaces, see [6, 7, 15, 17, 19, 25] in the reference section of this paper and others in the literature.

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

In this paper, we establish that it is possible to prove common coupled fixed point of mappings in the setting of partially ordered metric space without imposing the above outlined conditions on the mappings but by enhancing the initial assumptions of the theorem of Bhaskar and Lakshmikantam [1] together with the mixed monotone property. Our method is distinct from the existing ones in that we have enlarged the class of mappings that can be investigated due to the following reasons:
(i) We do not assume compatibility and weak compatibility of mappings involved;
(ii) We do not make one of the mappings a proper subset of the other;
(iii) We used our method to prove common coupled fixed point theorem for family of mappings; and
(iv) We do not impose comparison criterion a priori on the sequences involved in the computation of the common coupled fixed points.

2 Main Results

A. Results Involving Rational-type Contractive Conditions.

Theorem 2.1. Let $(X, \leq)$ be a partially ordered set and d a metric on X such that (X, d) is a complete metric space. Let $F, G : X \times X \to X$ be mappings having the mixed monotone property such that for some $\lambda \geq 0$, $\forall a, b, u, v \in X$, $d(u, F(u, v)) + d(a, u) > 0$ and ψ, a (c)-comparison function , we have

$$d(F(a,b), G(u,v)) \leq \frac{\lambda d(a, F(a,b)).d(a, G(u,v)).d(u, F(a,b))}{1 + d(u, F(u,v)) + d(a,u)} + \psi(d(a,u)). \tag{2.1}$$

Suppose that we endow the product space $X \times X$ with the following partial order:
For $(a,b), (u,v) \in X \times X$, $(u,v) \leq (a,b) \Longleftrightarrow a \geq u, b \leq v$.
If there exist $a_0, b_0 \in X$, such that $a_0 \leq F(a_0, b_0) \leq G(a_0, b_0)$ and $b_0 \geq F(b_0, a_0) \geq G(b_0, a_0)$. Then all F and G have a common coupled fixed point.

Proof. Choose $a_0, b_0 \in X$ such that $a_0 \leq F(a_0, b_0) \leq G(a_0, b_0)$ and $b_0 \geq F(b_0, a_0) \geq G(b_0, a_0)$.
Define $a_{2k+1} = F(a_{2k}, b_{2k})$, $b_{2k+1} = F(b_{2k}, a_{2k})$ and $a_{2k+2} = G(a_{2k+1}, b_{2k+1})$, $b_{2k+2} = G(b_{2k+1}, a_{2k+1})$ for $k \geq 0$.
We are to prove that a_k is non-decreasing and b_k is non-increasing. That is, for all $k \geq 0$,

$$a_{2k} \leq a_{2k+1} \leq a_{2k+2}$$

and

$$b_{2k} \geq b_{2k+1} \geq b_{2k+2}.$$

Firstly, $a_0 \leq F(a_0, b_0) = a_1$, and $b_0 \geq F(b_0, a_0) = b_1$.
By the iterative process above,

$$a_2 = G(a_1, b_1), \ b_2 = G(b_1, a_1).$$

Owing to the mixed monotone property of F and G, we have

$$a_1 = F(a_0, b_0) \leq G(a_0, b_0) \leq G(a_1, b_1) = a_2,$$

$$b_1 = F(b_0, a_0) \geq G(b_0, a_0) \geq G(b_1, a_1) = b_2,$$

Moreover,

$$a_2 = G(a_1, b_1) \leq G(a_2, b_2) = a_3,$$
$$b_2 = G(b_1, a_1) \geq G(b_2, a_2) = b_3$$

and

$$a_3 = F(a_2, b_2) \leq F(a_3, b_3) = a_4,$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

$$b_3 = F(b_2, a_2) \geq F(b_3, a_3) = b_4.$$

Therefore, for $n \geq 1$,

$$a_{2k+1} = F(a_{2k}, b_{2k}) \leq F(a_{2k+1}, b_{2k+1}) = a_{2k+2},$$
$$b_{2k+1} = F(b_{2k}, a_{2k}) \geq F(b_{2k+1}, a_{2k+1}) = b_{2k+2},$$

and

$$a_{2k+2} = G(a_{2k+1}, b_{2k+1}) \leq G(a_{2k+2}, b_{2k+2}) = a_{2k+3},$$
$$b_{2k+2} = G(b_{2k+1}, a_{2k+1}) \geq G(b_{2k+2}, a_{2k+2}) = b_{2k+3}.$$

Hence,

$$a_0 \leq a_1 \leq a_2 \leq ... \leq a_{2k} \leq a_{2k+1} \leq ...$$

and

$$b_0 \geq b_1 \geq b_2 \geq ... \geq b_{2k} \geq b_{2k+1} \geq ...$$

Therefore, we deduce by (2.1) that

$$\begin{aligned} d(a_{2k+1}, a_{2k+2}) &= d(F(a_{2k}, b_{2k}), G(a_{2k+1}, b_{2k+1})) \\ &\leq \frac{\lambda d(a_{2k}, F(a_{2k}, b_{2k})).d(a_{2k}, G(a_{2k+1}, b_{2k+1})).d(a_{2k+1}, F(a_{2k}, b_{2k}))}{1 + d(a_{2k+1}, F(a_{2k+1}, b_{2k+1})) + d(a_{2k}, a_{2k+1})} \\ &+ \psi(d(a_{2k}, a_{2k+1})) \\ &= \frac{\lambda d(a_{2k}, a_{2k+1}) d(a_{2k}, a_{2k+2}).d(a_{2k+1}, a_{2k+1})}{1 + d(a_{2k+1}, a_{2k+2}) + d(a_{2k}, a_{2k+1})} + \psi(d(a_{2k}, a_{2k+1})) \\ &= \psi(d(a_{2k}, a_{2k+1})). \end{aligned} \tag{2.2}$$

Similarly,

$$\begin{aligned} d(b_{2k+1}, b_{2k+2}) &= d(F(b_{2k}, a_{2k}), G(b_{2k+1}, a_{2k+1})) \\ &\leq \frac{\lambda d(b_{2k}, F(b_{2k}, a_{2k})).d(b_{2k}, G(b_{2k+1}, a_{2k+1})).d(b_{2k+1}, F(b_{2k}, a_{2k}))}{1 + d(b_{2k+1}, F(b_{2k+1}, a_{2k+1})) + d(b_{2k}, b_{2k+1})} \\ &+ \psi d(b_{2k}, b_{2k+1}) \\ &= \frac{\lambda d(b_{2k}, b_{2k+1}) d(b_{2k}, b_{2k+2}).d(b_{2k+1}, b_{2k+1})}{1 + d(b_{2k+1}, b_{2k+2}) + d(b_{2k}, b_{2k+1})} + \psi d(b_{2k}, b_{2k+1}) \\ &= \psi(d(b_{2k}, b_{2k+1})). \end{aligned} \tag{2.3}$$

Therefore,

$$d(a_{2k+1}, a_{2k+2}) \leq \psi(d(a_{2k}, a_{2k+1})) \tag{2.4}$$

and

$$d(b_{2k+1}, b_{2k+2}) \leq \psi(d(b_{2k}, b_{2k+1})) \tag{2.5}$$

Similarly, proceeding as above, we have

$$d(a_{2k+2}, a_{2k+3}) \leq \psi(d(a_{2k+1}, a_{2k+2})) \tag{2.6}$$

and

$$d(b_{2k+2}, b_{2k+3}) \leq \psi(d(b_{2k+1}, b_{2k+2})). \tag{2.7}$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Hence, it can be deduced from (2.4),(2.5), (2.6) and (2.7) that

$$\begin{aligned} d(a_{2k+2}, a_{2k+3}) + d(b_{2k+2}, b_{2k+3}) &\leq \psi(d(a_{2k+1}, a_{2k+2})) + \psi(d(b_{2k+1}, b_{2k+2})) \\ &\leq \psi^2(d(a_{2k}, a_{2k+1})) + \psi^2(d(b_{2k}, b_{2k+1})) \\ &\leq ... \leq \psi^n(d(a_0, a_1)) + \psi^n(d(b_0, b_1)). \end{aligned}$$

Thus, it follows that

$$d(a_{n+1}, a_n) + d(b_{n+1}, b_n) \leq \psi^n(d(a_0, a_1)) + \psi^n(d(b_0, b_1)). \tag{2.8}$$

Furthermore, for $n, r \in \mathbb{N}$, using (2.8) inductively and repeated application of triangle inequality, we have

$$\begin{aligned} d(a_n, a_{n+r}) + d(b_n, b_{n+r}) &\leq [d(a_n, a_{n+1}) + d(b_n, b_{n+1})] \\ &\quad + [d(a_{n+1}, a_{n+2}) + d(b_{n+1}, b_{n+2})] + ... \\ &\quad + [d(a_{n+r-1}, a_{n+r}) + d(b_{n+r-1}, b_{n+r})] \\ &\leq \psi^n(d(a_0, a_1)) + \psi^n(d(b_0, b_1)) + \psi^{n+1}(d(a_0, a_1)) \\ &\quad + \psi^{n+1}(d(b_0, b_1)) + ... + \psi^{n+r-1}(d(a_0, a_1)) + \psi^{n+r-1}(d(b_0, b_1)) \\ &= \psi^n(d(a_0, a_1)) + \psi^{n+1}(d(a_0, a_1)) + ... + \psi^{n+r-1}(d(a_0, a_1)) \\ &\quad + \psi^n(d(b_0, b_1)) + \psi^{n+1}(d(b_0, b_1)) + ... + \psi^{n+r-1}(d(b_0, b_1)) \end{aligned}$$

$$\begin{aligned} &= \sum_{k=n}^{n+r-1} \psi^k(d(a_0, a_1)) + \sum_{k=n}^{n+r-1} \psi^k(d(b_0, b_1)) \\ &= \sum_{k=0}^{n+r-1} \psi^k(d(a_0, a_1)) - \sum_{k=0}^{n-1} \psi^k(d(a_0, a_1)) + \sum_{k=0}^{n+r-1} \psi^k(d(b_0, b_1)) \\ &\quad - \sum_{k=0}^{n-1} \psi^k(d(b_0, b_1)) \quad \to 0 \ as \ n \to \infty \end{aligned}$$

since ψ is a (c)-comparison function. Therefore, $\{a_n\}$,$\{a_n\}$ are Cauchy sequences in (X, d). Furthermore, since (X, d) is a complete metric space, there exist $a^*, b^* \in X$ such that

$$\lim_{n\to\infty} a_n = a^*$$

and

$$\lim_{n\to\infty} b_n = b^*$$

.

We now show that (a^*, b^*) is a coupled fixed point of F and G.
By condition (2.1) again, and noting that $\psi(0) = 0$, we have

$$\begin{aligned} d(a^*, G(a^*, b^*)) &\leq d(a^*, a_{2n+1}) + d(a_{2n+1}, G(a^*, b^*)) \\ &= d(a^*, a_{2n+1}) + d(F(a_{2n}, b_{2n}), G(a^*, b^*)) \\ &\leq d(a^*, a_{2n+1}) + \frac{\lambda d(a_{2n}, F(a_{2n}, b_{2n})).d(a_{2n}, G(a^*, b^*)).d(a^*, F(a_{2n}, b_{2n}))}{1 + d(a^*, F(a^*, b^*)) + d(a^*, a_{2n})} \\ &\quad + \psi(d(a_{2n}, a^*)) \\ &= d(a^*, a_{2n+1}) + \frac{\lambda d(a_{2n}, a_{2n+1}) d(a_{2n}, G(a^*, b^*)).d(a^*, a_{2n+1}))}{1 + d(a^*, F(a^*, b^*)) + d(a_{2n}, a^*)} + \psi(d(a_{2n}, a^*)) \\ &\to 0 \ as \ n \to \infty. \end{aligned}$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Therefore $d(a^*, G(a^*, b^*)) = 0$, with the consequence that $a^* = G(a^*, b^*)$.
Again by (2.1), we have

$$\begin{aligned} d(b^*, G(b^*, a^*)) &\leq d(b^*, b_{2n+1}) + d(b_{2n+1}, G(b^*, a^*)) \\ &= d(b^*, b_{2n+1}) + d(F(b_{2n}, a_{2n}), G(a^*, b^*)) \\ &\leq d(b^*, b_{2n+1}) + \frac{\lambda d(b_{2n}, F(b_{2n}, a_{2n})).d(b_{2n}, G(b^*, a^*)).d(b^*, F(b_{2n}, a_{2n}))}{1 + d(b^*, F(b^*, a^*)) + d(b^*, b_{2n})} \\ &+ \psi(d(b_{2n}, b^*)) \\ &= d(b^*, b_{2n+1}) + \frac{\lambda d(b_{2n}, b_{2n+1}) d(b_{2n}, G(b^*, a^*)).d(b^*, b_{2n+1}))}{1 + d(b^*, F(b^*, a^*)) + d(b_{2n}, b^*)} + \psi(d(b_{2n}, b^*)) \\ &\to 0 \; as \; n \to \infty. \end{aligned}$$

Therefore $d(b^*, G(b^*, a^*)) = 0$, implying that $b^* = G(b^*, a^*)$.
Hence, (a^*, b^*) is a coupled fixed point of G.
Next is to show that (a^*, b^*) is a coupled fixed point of F too. By (2.1), we have

$$\begin{aligned} d(F(a^*, b^*), a^*) &= d(F(a^*, b^*), G(a^*, b^*)) \\ &\leq \frac{\lambda d(a^*, F(a^*, b^*)).d(a^*, G(a^*, b^*)).d(a^*, F(a^*, b^*))}{1 + d(a^*, F(a^*, b^*)) + d(a^*, a^*)} + \psi(d(a^*, a^*)) \\ &= \frac{\lambda d(a^*, F(a^*, b^*)).d(a^*, a^*).d(a^*, F(a^*, b^*))}{1 + d(a^*, F(a^*, b^*)) + d(a^*, a^*)} + \psi(0) = 0, \end{aligned}$$

implying that $d(F(a^*, b^*), a^*) = 0$, and by extension, $a^* = F(a^*, b^*)$.

Similarly by (2.1), we have

$$\begin{aligned} d(F(b^*, a^*), b^*) &= d(F(b^*, a^*), G(b^*, a^*)) \\ &\leq \frac{\lambda d(b^*, F(b^*, a^*)).d(b^*, G(b^*, a^*)).d(b^*, F(b^*, a^*))}{1 + d(b*, F(b^*, a^*)) + d(b^*, b^*)} + \psi(d(b^*, b^*)) \\ &= \frac{\lambda d(b^*, F(b^*, a^*)) d(b^*, b^*).d(b^*, F(b^*, a^*))}{1 + d(b^*, F(b^*, a^*)) + d(b^*, b^*)} + \psi(0) = 0, \end{aligned}$$

implying that $d(F(b^*, a^*), b^*) = 0$, therefore, $b^* = F(b^*, a^*)$.
Thus, (a^*, b^*) is a common coupled fixed point of F and G.
Example 2.2. Let $X = [0, 4]$ be endowed with the usual metric. Let $\psi(t) = \frac{1}{2}t$, for all $t \in X$. Clearly, $\psi(t)$ is a (c)-comparison function. Define $F, G : [0, 4] \times [0, 4] \to [0, 4]$ by:

$$F(a, b) = 3a - 2b,$$

$$G(a, b) = \begin{cases} 2a - b + 1, & \text{if } a \geq b; \\ \frac{1}{7}, & \text{if } a < b. \end{cases}$$

Clearly, F and G have the mixed monotone property.
Let $a_0 = \frac{2}{3}$, $b_0 = \frac{1}{2} \in X$.
$F(a_0, b_0) = F(\frac{2}{3}, \frac{1}{2}) = 3(\frac{2}{3}) - 2(\frac{1}{2}) = 1$
$F(b_0, a_0) = F(\frac{1}{2}, \frac{2}{3}) = 3(\frac{1}{2}) - 2(\frac{2}{3}) = \frac{1}{6}$.
$G(a_0, b_0) = G(\frac{2}{3}, \frac{1}{2}) = 2(\frac{2}{3}) - (\frac{1}{2}) + 1 = \frac{11}{6}$
$F(b_0, a_0) = G(\frac{1}{2}, \frac{2}{3}) = \frac{1}{7}$.
Thus,
$\frac{2}{3} < 1 < \frac{11}{6}$ and $\frac{1}{2} > \frac{1}{6} > \frac{1}{7}$.
Hence,
$a_0 \leq F(a_0, b_0) \leq G(a_0, b_0)$ and $b_0 \geq F(b_0, a_0) \geq G(b_0, a_0)$.
Consider the contractive condition (2.1) in Theorem 2.1. Let $a = \frac{4}{3}$, $b = \frac{1}{3}$, $\lambda = 2$, $u = \frac{4}{5}$ and $v = \frac{1}{5}$.

IJMSO
International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Then, $F(a,b) = F(\frac{4}{3}, \frac{1}{3}) = \frac{10}{3}$, $G(u,v) = G(\frac{4}{5}, \frac{1}{5}) = \frac{12}{5}$, $d(F(a,b), G(u,v)) = d(F(\frac{4}{3}, \frac{1}{3}), G(\frac{4}{5}, \frac{1}{5})) = \left|\frac{10}{3} - \frac{12}{5}\right| = \frac{14}{15}$, $F(u,v) = 3(\frac{4}{5}) - 2(\frac{1}{5}) = 2$, $d(a, F(a,b)) = \left|\frac{4}{3} - \frac{10}{3}\right| = 2$, $d(a, G(u,v)) = \left|\frac{4}{3} - \frac{12}{5}\right| = \frac{16}{15}$, $d(u, F(a,b)) = \left|\frac{4}{5} - \frac{10}{3}\right| = \frac{38}{15}$, $d(a,u) = \left|\frac{4}{3} - \frac{4}{5}\right| = \frac{8}{15}$ and $d(u, F(u,v)) = \left|\frac{4}{5} - 2\right| = \frac{6}{5}$.
Hence, contractive condition (2.1), i.e

$$d(F(a,b), G(u,v)) \leq \frac{\lambda d(a, F(a,b)).d(a, G(u,v)).d(u, F(a,b))}{1 + d(u, F(u,v)) + d(a,u)} + \psi(d(a,u)), \tag{2.9}$$

implies that $\frac{14}{15} = 0.9333 \leq \frac{2\times 2\times \frac{16}{15}\times\frac{38}{15}}{1+\frac{6}{5}+\frac{8}{15}} + \frac{1}{2}(\frac{8}{15}) = \frac{2432}{615} + \frac{8}{30} = 4.2211$.
Since all the assumptions of Theorem 2.1 are satisfied, F and G have a common coupled fixed point in $X = [0,4]$.

Corollary 2.3. Let $(X, \leq)$ be a partially ordered set and d a metric on X such that (X,d) is a complete metric space. Let $F, G : X \times X \to X$ be mappings having the mixed monotone property such that for some $\lambda \geq 0$, $\mu \in [0,1)$ and $\forall a, b, u, v \in X$, where $d(u, F(u,v)) + d(a,u) > 0$, we have

$$d(F(a,b), G(u,v)) \leq \frac{\lambda d(a, F(a,b)).d(a, G(u,v)).d(u, F(a,b))}{1 + d(u, F(u,v)) + d(a,u)} + \mu d(a,u). \tag{2.10}$$

Suppose that we endow the product space $X \times X$ with the following partial order:
For $(a,b), (u,v) \in X \times X, (u,v) \leq (a,b) \iff a \geq u, b \leq v$.
If there exist $a_0, b_0 \in X$, such that $a_0 \leq F(a_0, b_0) \leq G(a_0, b_0)$ and $b_0 \geq F(b_0, a_0) \geq G(b_0, a_0)$.
Then all F and G have a common coupled fixed point.

B. Results Involving Contractive Conditions of Non-Rational-Type.

Theorem 2.4. Let (X,d) be a complete metric space and $(X, \leq)$ a partially ordered set. Suppose $\{T_\alpha\}_{\alpha \in J}$ with $T_\alpha : X \times X \to X$ is a family of mappings and J is an index set. If there exists a fixed $\beta \in J$ such that for each $\alpha \in J$,

$$\begin{aligned} d(T_\alpha(s,r), T_\beta(u,v)) \leq\ & a d(s,u) + b[d(s, T_\alpha(s,r)) + d(u, T_\beta(u,v))] \\ & + c[d(s, T_\beta(u,v)) + d(u, T_\alpha(s,r))] \end{aligned} \tag{2.11}$$

$\forall s, r, u, v \in X$, for some $a, b, c \geq 0$ and $a + 2b + 2c < 1$.
Suppose that we endow the product space $X \times X$ with the following partial order:
For $(s,r), (u,v) \in X \times X, (u,v) \leq (s,r) \iff s \geq u, r \leq v$.
If $\forall \alpha \in J$, there exist $s_0, r_0 \in X$, such that $s_0 \leq T_\alpha(s_0, r_0) \leq T_\beta(s_0, r_0)$ and $r_0 \geq T_\alpha(r_0, s_0) \geq T_\beta(r_0, s_0)$. Then all T_α have a common coupled fixed point.

Proof. Choose $s_0, r_0 \in X$ such that $s_0 \leq T_\alpha(s_0, r_0) \leq T_\beta(s_0, r_0)$ and $r_0 \geq T_\alpha(r_0, s_0) \geq T_\beta(r_0, s_0)$. and define sequences $\{s_n\}$, $\{r_n\}$ in X in the following way:

$$s_{2n+1} = T_\alpha(s_{2n}, r_{2n}),\ r_{2n+1} = T_\alpha(r_{2n}, s_{2n})$$

$$s_{2n+2} = T_\beta(s_{2n+1}, r_{2n+1}),\ r_{2n+2} = T_\beta(r_{2n+1}, s_{2n+1})$$

We are to prove that $\{s_n\}$ is non-decreasing and $\{r_n\}$ is non-increasing. That is, for all $n \geq 0$, $s_{2n} \leq s_{2n+1}$ and $r_{2n} \geq r_{2n+1}$
Firstly, we set

$$s_0 \leq T_\alpha(s_0, r_0) = s_1,$$

(say) and

$$r_0 \geq T_\alpha(r_0, s_0) = r_1$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

(say).
By the iterative process above,

$$s_2 = T_\beta(s_1, r_1),\ r_2 = T_\beta(r_1, s_1).$$

Owing to the mixed monotone property of T_α and T_β, we have

$$s_1 = T_\alpha(s_0, r_0) \leq T_\beta(s_0, r_0) \leq T_\beta(s_1, r_1) = s_2,$$

$$r_1 = T_\alpha(r_0, s_0) \geq T_\beta(r_0, s_0) \geq T_\beta(r_1, s_1) = r_2.$$

Moreover,

$$s_2 = T_\beta(s_1, r_1) \leq T_\beta(s_2, r_2) = s_3,$$
$$r_2 = T_\beta(r_1, s_1) \geq T_\beta(r_2, s_2) = r_3,$$

and

$$s_3 = T_\alpha(s_2, r_2) \leq T_\alpha(s_3, r_3) = s_4,$$
$$r_3 = T_\alpha(r_2, s_2) \geq T_\alpha(r_3, s_3) = r_4.$$

Therefore, for $n \geq 1$,

$$s_{2n+1} = T_\alpha(s_{2n}, r_{2n}) \leq T_\alpha(s_{2n+1}, r_{2n+1}) = s_{2n+2},$$
$$r_{2n+1} = T_\alpha(r_{2n}, s_{2n}) \geq T_\alpha(r_{2n+1}, s_{2n+1}) = r_{2n+2},$$

and

$$s_{2n+2} = T_\beta(s_{2n+1}, r_{2n+1}) \leq T_\beta(s_{2n+2}, r_{2n+2}) = s_{2n+3},$$
$$r_{2n+2} = T_\beta(r_{2n+1}, s_{2n+1}) \geq T_\beta(r_{2n+2}, s_{2n+2}) = r_{2n+3}.$$

Hence,

$$s_0 \leq s_1 \leq s_2 \leq ... \leq s_{2n} \leq s_{2n+1} \leq ...,$$
$$r_0 \geq r_1 \geq r_2 \geq ... \geq r_{2n} \geq r_{2n+1} \geq ....$$

Therefore, we deduce by (2.11) that

$$\begin{aligned}
d(s_{2n+1}, s_{2n+2}) &= d(T_\alpha(s_{2n}, r_{2n}), T_\beta(s_{2n+1}, r_{2n+1})) \\
&\leq ad(s_{2n}, s_{2n+1}) + b[d(s_{2n}, T_\alpha(s_{2n}, r_{2n})) + d(s_{2n+1}, T_\beta(s_{2n+1}, r_{2n+1}))] \\
&+ c[d(s_{2n}, T_\beta(s_{2n+1}, r_{2n+1})) + d(s_{2n+1}, T_\alpha(s_{2n}, r_{2n}))] \\
&\leq ad(s_{2n}, s_{2n+1}) + b[d(s_{2n}, s_{2n+1}) + d(s_{2n+1}, s_{2n+2})] \\
&+ c[d(s_{2n}, s_{2n+2}) + d(s_{2n+1}, s_{2n+1})] \\
&= ad(s_{2n}, s_{2n+1}) + bd(s_{2n}, s_{2n+1}) + bd(s_{2n+1}, s_{2n+2}) + cd(s_{2n}, s_{2n+2}) \qquad (2.12) \\
&\leq ad(s_{2n}, s_{2n+1}) + bd(s_{2n}, s_{2n+1}) + bd(s_{2n+1}, s_{2n+2}) + cd(s_{2n}, s_{2n+1}) \\
&+ cd(s_{2n+1}, s_{2n+2}) \\
&\leq \frac{a+b+c}{1-b-c} d(s_{2n}, s_{2n+1}) \\
&= qd(s_{2n}, s_{2n+1}),
\end{aligned}$$

where

$$q = \frac{a+b+c}{1-b-c} < 1.$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Similarly by (2.11),

$$\begin{aligned} d(r_{2n+1}, r_{2n+2}) &= d(T_\alpha(r_{2n}, s_{2n}), T_\beta(r_{2n+1}, s_{2n+1})) \\ &\leq ad(r_{2n}, r_{2n+1}) + b[d(r_{2n}, T_\alpha(r_{2n}, s_{2n})) + d(r_{2n+1}, T_\beta(r_{2n+1}, s_{2n+1}))] \\ &+ c[d(r_{2n}, T_\beta(r_{2n+1}, s_{2n+1})) + d(r_{2n+1}, T_\alpha(r_{2n}, s_{2n}))] \\ &\leq ad(r_{2n}, r_{2n+1}) + b[d(r_{2n}, r_{2n+1}) + d(r_{2n+1}, r_{2n+2})] \\ &+ c[d(r_{2n}, r_{2n+2}) + d(r_{2n+1}, r_{2n+1})] \\ &= ad(r_{2n}, r_{2n+1}) + bd(r_{2n}, r_{2n+1}) + bd(r_{2n+1}, r_{2n+2}) + cd(r_{2n}, r_{2n+2}) \\ &\leq ad(r_{2n}, r_{2n+1}) + bd(r_{2n}, r_{2n+1}) + bd(r_{2n+1}, r_{2n+2}) + cd(r_{2n}, r_{2n+1}) \\ &+ cd(r_{2n+1}, r_{2n+2}) \\ &\leq ad(r_{2n}, r_{2n+1}) + bd(r_{2n}, r_{2n+1}) + bd(r_{2n+1}, r_{2n+2}) + c[d(r_{2n}, r_{2n+1}) \\ &+ d(r_{2n+1}, r_{2n+2})] \\ &\leq \frac{a+b+c}{1-b-c} d(r_{2n}, r_{2n+1}) = qd(r_{2n}, r_{2n+1}). \end{aligned} \tag{2.13}$$

Now, adding (2.12) and (2.13), we have

$$d(s_{2n+1}, s_{2n+2}) + d(r_{2n+1}, r_{2n+2}) \leq q[d(s_{2n}, s_{2n+1}) + d(r_{2n}, r_{2n+1})]$$

Similarly by (2.11),

$$\begin{aligned} d(s_{2n}, s_{2n+1}) &= d(T_\alpha(s_{2n-1}, r_{2n-1}), T_\beta(s_{2n}, r_{2n})) \\ &\leq ad(s_{2n-1}, s_{2n}) + b[d(s_{2n-1}, T_\alpha(s_{2n-1}, r_{2n-1})) + d(s_{2n}, T_\beta(s_{2n}, r_{2n}))] \\ &+ c[d(s_{2n-1}, T_\beta(s_{2n}, r_{2n})) + d(s_{2n}, T_\alpha(s_{2n-1}, r_{2n-1}))] \\ &\leq ad(s_{2n-1}, s_{2n}) + b[d(s_{2n-1}, s_{2n}) + d(s_{2n}, s_{2n+1})] \\ &+ c[d(s_{2n-1}, s_{2n+1}) + d(s_{2n}, s_{2n})] \\ &\leq ad(s_{2n-1}, s_{2n}) + bd(s_{2n-1}, s_{2n}) + bd(s_{2n}, s_{2n+1}) + cd(s_{2n-1}, s_{2n}) \\ &+ cd(s_{2n}, s_{2n+1}) \\ &\leq \frac{a+b+c}{1-b-c} d(s_{2n-1}, s_{2n}) = qd(s_{2n-1}, s_{2n}). \end{aligned} \tag{2.14}$$

Furthermore, by (2.11), it follows that

$$\begin{aligned} d(r_{2n}, r_{2n+1}) &= d(T_\alpha(r_{2n-1}, s_{2n-1}), T_\beta(r_{2n}, s_{2n})) \\ &\leq ad(r_{2n}, r_{2n-1}) + b[d(r_{2n-1}, T_\alpha(r_{2n-1}, s_{2n-1})) + d(r_{2n}, T_\beta(r_{2n}, s_{2n}))] \\ &+ c[d(r_{2n-1}, T_\beta(r_{2n}, s_{2n})) + d(r_{2n}, T_\alpha(r_{2n-1}, s_{2n-1}))] \\ &\leq ad(r_{2n-1}, r_{2n}) + b[d(r_{2n-1}, r_{2n}) + d(r_{2n}, r_{2n+1})] \\ &+ c[d(r_{2n-1}, r_{2n+1}) + d(r_{2n}, r_{2n})] \\ &\leq ad(r_{2n-1}, r_{2n}) + bd(r_{2n-1}, r_{2n}) + bd(r_{2n}, r_{2n+1}) + c[d(r_{2n-1}, r_{2n+1}) \\ &\leq \frac{a+b+c}{1-b-c} d(r_{2n}, r_{2n+1}) = qd(r_{2n-1}, r_{2n}). \end{aligned} \tag{2.15}$$

Also, from (2.14) and (2.15), we have

$$d(s_{2n}, s_{2n+1}) + d(r_{2n}, r_{2n+1}) \leq q[d(s_{2n-1}, s_{2n}) + d(r_{2n-1}, r_{2n})]. \tag{2.16}$$

If we let

$$\varepsilon_n = d(s_{2n}, s_{2n+1}) + d(r_{2n}, r_{2n+1}),$$

then by (2.16), we have

$$\varepsilon_n \leq q\varepsilon_{n-1}. \tag{2.17}$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

Thus,

$$0 \leq \varepsilon_n \leq q\varepsilon_{n-1} \leq q^2\varepsilon_{n-2} \leq ... \leq q^n\varepsilon_0. \tag{2.18}$$

If $\varepsilon_0 = 0$, then (s_0, r_0) is a coupled fixed point of T. Suppose that $\varepsilon_0 > 0$. Then for each $m \in \mathbb{N}$, we obtain by repeated application of triangle inequality and (2.18) that

$$\begin{aligned} d(s_{2n}, s_{2n+m}) + d(r_{2n}, r_{2n+m}) &\leq [d(s_{2n}, s_{2n+1}) + d(r_{2n}, r_{2n+1})] + \\ &\quad [d(s_{2n+1}, s_{2n+2}) + d(r_{2n+1}, r_{2n+2})] + ... \\ &\quad + [d(s_{2n+m-1}, s_{2n+m}) + d(r_{2n+m-1}, r_{2n+m})] \\ &= \varepsilon_n + \varepsilon_{n+1} + ... + \varepsilon_{n+m-1} \\ &\leq q^n\varepsilon_0 + q^{n+1}\varepsilon_0 + ... + q^{n+m-1}\varepsilon_0 \\ &\leq q^n\varepsilon_0 + q^{n+1}\varepsilon_0 + ... + q^{n+m-1}\varepsilon_0 \\ &= \frac{q^n(1-q^{m-1})\varepsilon_0}{1-q} \to 0 \ as \ n \to \infty. \end{aligned} \tag{2.19}$$

Therefore, $\{s_n\}$,$\{r_n\}$ are Cauchy sequences in (X, d). Since (X, d) is a complete metric space, there exist $s^*, r^* \in X$ such that $\lim_{n\to\infty} s_n = s^*$ and $\lim_{n\to\infty} r_n = r^*$.
We first show that (s^*, r^*) is a coupled fixed point of T_β. By condition (2.11) again, we have

$$\begin{aligned} d(s^*, T_\beta(s^*, r^*)) &\leq d(s^*, s_{2n+1}) + d(s_{2n+1}, T_\beta(s^*, r^*)) \\ &= d(s^*, s_{2n+1}) + d(T_\alpha(s_{2n}, r_{2n}), T_\beta(s^*, r^*)) \\ &\leq d(s^*, s_{2n+1}) + ad(s_{2n}, s^*) + b[d(s_{2n}, T_\alpha(s_{2n}, r_{2n})) + d(s^*, T_\beta(s^*, r^*))] \\ &+ c[d(s_{2n}, T_\beta(s^*, r^*)) + d(s^*, T_\alpha(s_{2n}, r_{2n}))] \\ &= d(s^*, s_{2n+1}) + ad(s_{2n}, s^*) + b[d(s_{2n}, s_{2n+1}) + bd(s^*, T_\beta(s^*, r^*))] \\ &+ c[d(s_{2n}, T_\beta(s^*, r^*)) + d(s^*, s_{2n+1})] \\ &\leq d(s^*, s_{2n+1}) + ad(s_{2n}, s^*) + b[d(s_{2n}, s^*) + d(s^*, s_{2n+1}) + bd(s^*, T_\beta(s^*, r^*))] \\ &+ c[d(s_{2n}, s^*) + d(s^*, T_\beta(s^*, r^*)) + d(s^*, s_{2n+1})] \\ &\leq \frac{d(s^*, s_{2n+1}) + ad(s_{2n}, s^*) + b[d(s_{2n}, s^*) + d(s^*, s_{2n+1})]}{1-b-c} \\ &+ \frac{c[d(s_{2n}, s^*) + d(s^*, s_{2n+1})]}{1-b-c} \\ &\to 0 \ \ as \ n \to \infty. \end{aligned} \tag{2.20}$$

Therefore $d(s^*, T_\beta(s^*, r^*)) = 0$, implying that $s^* = T_\beta(s^*, r^*)$.
Similarly,

$$\begin{aligned} d(r^*, T_\beta(r^*, s^*)) &\leq d(r^*, r_{2n+1}) + d(r_{2n+1}, T_\beta(r^*, s^*)) \\ &= d(r^*, r_{2n+1}) + d(T_\alpha(r_{2n}, s_{2n}), T_\beta(r^*, s^*)) \\ &\leq d(r^*, r_{2n+1}) + ad(r_{2n}, r^*) + b[d(r_{2n}, T_\alpha(r_{2n}, s_{2n})) + d(r^*, T_\beta(r^*, s^*))] \\ &+ c[d(r_{2n}, T_\beta(r^*, s^*)) + d(r^*, T_\alpha(r_{2n}, s_{2n}))] \\ &= d(r^*, r_{2n+1}) + ad(r_{2n}, r^*) + b[d(r_{2n}, r_{2n+1}) + bd(r^*, T_\beta(r^*, s^*))] \\ &+ c[d(r_{2n}, T_\beta(r^*, s^*)) + d(r^*, r_{2n+1})] \\ &\leq d(r^*, r_{2n+1}) + ad(r_{2n}, r^*) + b[d(r_{2n}, r^*) + d(r^*, r_{2n+1}) + d(r^*, T_\beta(r^*, s^*))] \\ &+ c[d(r_{2n}, r^*) + d(r^*, T_\beta(r^*, s^*)) + d(r^*, r_{2n+1})] \\ &\leq \frac{d(r^*, r_{2n+1}) + ad(r_{2n}, r^*) + b[d(r_{2n}, r^*) + d(r^*, r_{2n+1})]}{1-b-c} \\ &+ \frac{c[d(r_{2n}, r^*) + d(r^*, r_{2n+1})]}{1-b-c} \\ &\to 0 \ \ as \ n \to \infty, \end{aligned}$$

IJMSO

International Journal of Mathematical Sciences and Optimization: Theory and Applications
Vol. 8, No. 1, pp. 88 - 100
https://doi.org/10.6084/m9.figshare.20716042

implying that

$$d(r^*, T_\beta(r^*, s^*)) = 0$$

and thus

$$r^* = T_\beta(r^*, s^*).$$

Therefore, (s^*, r^*) is a coupled fixed point of T_β.
We now show that (s^*, r^*) is a coupled fixed point of $\{T_\alpha\}$ too.
Let $\alpha \in J$ be arbitrary, then by (2.11), we have

$$\begin{aligned} d(T_\alpha(s^*, r^*), s^*) &= d(T_\alpha(s^*, r^*), T_\beta(s^*, r^*)) \\ &\leq ad(s^*, s^*) + bd(s^*, T_\alpha(s^*, r^*)) + b[d(s^*, T_\beta(s^*, r^*)) \\ &+ cd(s^*, T_\beta(s^*, r^*))] + cd(s^*, T_\alpha(s^*, r^*)) \\ &= bd(s^*, T_\alpha(s^*, r^*)) + cd(s^*, T_\alpha(s^*, r^*)) \\ &= (b + c)d(s^*, T_\alpha(s^*, r^*)). \end{aligned}$$

Since $b + c < 1$, we get $d(s^*, T_\alpha(s^*, r^*)) = 0$, that is, $s^* = T_\alpha(s^*, r^*)$.
On further employing (2.11), it follows that

$$\begin{aligned} d(T_\alpha(r^*, s^*), r^*) &= d(T_\alpha(r^*, s^*), T_\beta(r^*, s^*)) \\ &\leq ad(r^*, r^*) + bd(r^*, T_\alpha(r^*, s^*)) + bd(r^*, T_\beta(r^*, s^*)) \\ &+ cd(r^*, T_\beta(r^*, s^*)) + cd(r^*, T_\alpha(r^*, s^*)) \\ &= bd(r^*, T_\alpha(r^*, s^*)) + cd(r^*, T_\alpha(r^*, s^*)) \\ &= (b + c)d(r^*, T_\alpha(r^*, s^*)). \end{aligned}$$

Since $(b + c) < 1$, we get $d(r^*, T_\alpha(r^*, s^*)) = 0$, that is $r^* = T_\alpha(r^*, s^*)$.
Thus, (s^*, r^*) is a coupled fixed point of T_α too.
Hence, (s^*, r^*) is a common coupled fixed point of T_α and T_β.

Remark 2.5. Theorem 2.5 is a variant of Theorem 2.1 of Abbas *et − al* [5] to the common coupled fixed point setting, involving family of mappings.
Remark 2.6. Furthermore, Theorem 2.5 generalizes and extends Theorems 2.5 and 2.6 of Sabetghadem *et al*. [23], to the common coupled fixed point setting in partially ordered space, the latter consisting of coupled fixed point in cone metric space setting.

3 Competing Interests

The authors of this paper declared that there are no competing interests as regards this article.

4 Acknowlegement

The authors wish to thank the reviewers for their helpful suggestions which further improve this research work.

References

[1] T. G. Bhaskar and V. Lakshmikantham; *Fixed point theorems in partially ordered metric spaces and applications*, Nonlinear Analysis: Theory, Methods and Application, 65 (7)(2006), 1379 - 1393.

[2] M. O. Olatinwo and K. R. Tijani; *Some results on stability of coupled fixed point iterations in metric spaces*, J. Adv. Math, Stud.,**13**(2)(2020), 169 - 178.

[3] K. R. Tijani and M. O. Olatinwo; *Some coupled fixed point theorems for mappings satisfying rational type contractive conditions in partially ordered metric spaces*, JP Journal of Fixed Point Theory and Applications,Volume 15, Number 1(2020), 19-43.

[4] M. Abbas, D. Ilic, and A. M. Khan; *Coupled coincidence point and common fixed point theorems in Partially ordered metric spaces with w-distance*, Fixed Point Theory and Applications, Volume 2010, article ID 134897, (2010) 11 pages.

[5] M. Abbas,T. Nazir, S. Radenovic; *Common coupled fixed points of generalized contractive mappings in partially ordered metric spaces*, Positivity, Doi 10.1007/s11117-012-0219-2,(2012)

[6] A. Aghajani, M. Abbas and P. Kallehbasti; *Common coupled fixed point theorems in partially ordered metric spaces and applications*, Mathematical Communications (2012).

[7] A. Aliouche; *Common fixed point theorems of Gregus type for weakly compatible mappings satisfying generalized contractive conditions*,Journal of mathematical Analysis and Applications, 341 (2008), 707 - 719.

[8] I. Beg, A. Latif, R. Ali and A. Azam; *Coupled fixed point of mixed monotone operators on probabilistic Banach spaces* , Archivum math 37(1)(2001), 1 - 8.

[9] Berinde, V; *Iterative approximation of fixed points*, Editura Efemeride (2002).

[10] V. Berinde, V; *Iterative approximation of fixed points*, Volume (1912) of Lecture Notes in Mathematics, Springer, Berlin, Second edition (2007)

[11] V. Berinde; *Generalized coupled fixed point theorems for mixed monotone mappings in partially ordered metric spaces*, Maths. F.A.,ar Xiv: 1103 - 5285vi (2012).

[12] N. Bilgili, I, M. Erhan, E. Karapinar and D. Turkoglu; *A note on 'coupled fixed point theorems for mixed g-monotone mappings in partially ordered metric spaces*, Fixed Point Theory and Applications, 2014, (2014) : 120.

[13] S. S. Chang and Y. H. Ma; *Coupled Fixed Points of mixed monotone Condensing operators and existence theorem of the solution for a class of functional equations arising in dynamic programming*, J. Math. Annal. Appl., 160(1991) 468 - 479.

[14] L. Ciric, M. O. Olatinwo, D. Gopal and G. Akinbo; *Coupled random fixed point theorems for mappings satisfying a contractive condition of rational type on a Partially Ordered Metric Space*, Advances in Fixed Point Theory, 2(1)(2012), 1 - 8.

[15] M. E. Gordji, E. Akbartabar, Y. J. Cho and M. Ramezani; *Coupled common fixed point theorems for mixed weakly monotone mappings in partially ordered metric spaces*, Fixed Point Theory and Applications, 2012, (2012) : 95.

[16] Z. Kadelburg, P, Kumam, S. Radenovic and W. Situnavarat; *Common coupled fixed point theorems for Geraghty-type contraction mappings using monotone property*, Fixed Point Theory and Applications,2015, (2015):27.

[17] J. K. Kim and S. Chandok; *Coupled common fixed point theorems for generalized nonlinear contraction mappings with mixed monotone property in partially ordered metric spaces*, Fixed Point Theory and Applications, 2012, (2012):95.

[18] V. Lakshmikantham and L. Ciric; *Coupled fixed point theorems for nonlinear contractions in partially ordered metric spaces*, Nonlinear Analysis: Theory, Methods and Applications, 70 (12)(2009), 4341 - 4349.

[19] A. Mutlu, N. Mermin, B. Mutlu and N. Bildik, ;*On common coupled fixed point theorems for comparable mappings in ordered partial metric spaces*, Acta Universitatis Apulensis, 23(2014), 91 - 98.

[20] M. O. Olatinwo;*Coupled fixed point theorems in cone metric spaces*, Ann. Univ. Ferrara ,**57**(1)(2011), 71 - 83.

[21] M. O. Olatinwo; *Coupled common fixed points of contractive mappings in metric spaces*, Journal of Advanced Research in Pure Mathematics, **4** (2)(2012), 11 - 20.

[22] M. O. Olatinwo and K. R. Tijani; *Some stability results for coupled fixed point iterative process in a complete metric spaces*, Surveys in Mathematics and its Applications,**14**(2019), 327 - 339.

[23] F. Sabetghadam, H. P. Masiha and A. H. Sanatpour; *Some coupled fixed points theorems in cone metric spaces*, Fixed Point Theory and Applications, Volume 2009, Article ID 125426, 8 pages, DoI:10.1155/2009/125426,(2009).

[24] M. Sarwar, S. Hussain and P. S. F. Kumati; *Common coupled fixed points theorems satisfying rational type contractive conditions in b-metric spaces*, SpringerPlus, (2016)5:257 DOI 10.1186/s40064-016-1849-6 ,(2016).

[25] W. Sintunavarat and P. Kumam; *Coupled coincidence and coupled common fixed point theorems in Partially Ordered Metric Spaces*, Thai Journal of Mathematics , 10(3) (2012), 551 - 563.

[26] C. A. Summit; *Some common fixed point results for rational type contraction mappings in partially ordered metric spaces*, Mathematical Bohemica, 138(4)(2013), 407 - 413.

[27] C. Summit, D. N. Tulsi, and M. A. Taoudi; *Fixed point theorem for generalized contractions satisfying rational type expressions in partially ordered metric spaces*, Gulf Journal of Mathematics, 2(4)(2014), 87 - 93.

[28] C. Summit, D. N. Tulsi and M. A. Taoudi; *Some coupled fixed point theorems for mappings satisfying a generalized contractive condition of rational type*, Palestine Journal of Mathematics, 4(2)(2015), 360 - 366.