Journal of the Nigerian Association of Mathematical Physics
Volume 30, (May, 2015), pp463 – 466
© J. of NAMP

Computational Error Estimate for the Power Series Solution of Odes Using Zeros of Chebyshev Polynomial

Issa K.[1], Ibrahim G.R.[2] and Bakare G.N.[3]

[1,2]Department of Statistics and Mathematical Sciences, Kwara State University, Malete, Nigeria.
[3]Department of Mathematics, University of Ilorin, Ilorin Nigeria.

Abstract

This paper compares the error estimation of power series solution with recursive Tau method for solving ordinary differential equations. From the computational viewpoint, the power series using zeros of Chebyshevpolunomial is effective, accurate and easy to use.

Keywords: Lanczos Tau method, Chebyshev polynomial, initial value problems, Lanczos – Ortiz Canonical polynomial, Ordinary Differential Equations

1.0 Introduction

This paper is concerned with the error estimated of ordinary differential equation of the form:

$$Ly(x) \equiv \sum_{r=0}^{m} \left(\sum_{k=0}^{N_r} P_{rk} x^k \right) y^{(r)}(x) = \sum_{r=0}^{\sigma} f_r x^r, \; a \le x \le b \tag{1}$$

together with the associated conditions:

$$L^* y(x_{rk}) \equiv \sum_{r=0}^{m-1} a_{rk} y^{(r)}(x_{rk}) = \alpha_k, \; k = 1(1)m \tag{2}$$

by seeking an approximate solution of the form

$$y_n(x) = \sum_{r=0}^{n} a_r x^r, \; r < +\infty$$

of $y(x)$ which is the exact solution of the corresponding perturbed system

$$Ly_n(x) = \sum_{r=0}^{\sigma} f_r x^r + H_n(x) \tag{3}$$

$$L^* y_n(x_r k) = \alpha_k, k = 1(1)m \tag{4}$$

where L is the linear differential operator, and α_k, f_r, $P_{r,k}$, N_r, $r = 0(1)m$, $k = 0(1)N_r$, a and b are real constants, $y^{(r)}$ denotes the derivatives of order r of $y(x)$, and the perturbation term $H_n(x)$ in equation (3) is defined by:

$$H_n(x) = \sum_{i=0}^{m+s-1} \tau_{i+1} T_{n-m+i+1}(x) = \sum_{i=0}^{m+s-1} \tau_{i+1} \sum_{i=0}^{n-m+i+1} C_r^{(n-m+i+1)} x^r \tag{5}$$

and $C_r^{(n)}$ is the coefficient of x^r in the $n-th$ degree chebyshev polynomial $T_n(x)$: that is,

$$T_n(x) = \cos\left(n \cos^{-1}\left\{ \frac{2x-a-b}{b-a} \right\} \right) \equiv \sum_{r=0}^{n} C_r^{(n)} x^r \tag{6}$$

Corresponding author: Issa K., E-mail: issakazeem@yahoo.com, Tel.: +2348036554437

The $\tau's$ are free parameters to be determined and s, the number of over determination of (1), is defined in [1-10]:

$$s = \max\{N_r - r > 0 | 0 \le r \le m\} \tag{7}$$

2.0 Review of Recursive formulation of the Tau approximant

In this section, we review the recursive approximant [2, 4, 11, 12, 13] and canonical polynomial [14], by adding perturbation terms to the right hand sides of (1). We have in [4]:

$$y_n(x) = \sum_{r=s}^{\sigma} f_r q_r(x) + \sum_{i=0}^{m+s-1} \tau_{i+1} \sum_{r=s}^{n-m+i+1} C_r^{(n-m+i+1)} q_r(x) \tag{8}$$

Assume $Q_r(x) = P_r = 1, r = 0(1)(s-1)$, then

$$\sum_{i=0}^{m+s-1} \tau_{i+1} \sum_{r=0}^{n-m+i+1} C_r^{(n-m+i+1)} P_r + \sum_{r=0}^{\sigma} f_r P_r = 0 \tag{9}$$

Where $q_r(x) = Q_r(x) - P_r$ and for the undetermined canonical polynomials (if any) assume $P_r = 1$, when equating the coefficient of $Q_r(x)$ to zero, otherwise $P_r = 0$ for $r = 0, 1, \cdots, (s-1)$ (that is equation (9) is the coefficient of undetermined canonical polynomials),

$$P_r = \frac{-1}{\sum_{k=0}^{m} k!\binom{r-s}{k} P_{k,k+s}} \left\{ \sum_{k=1}^{m} \left(\sum_{j=k}^{m} j!\binom{r-s}{j} P_{j,j-k} \right) P_{r-s-k} \right. \tag{10}$$

$$\left. + \sum_{k=0}^{s-1} \left(\sum_{j=0}^{m} j!\binom{r-s}{j} P_{j,j+k} \right) P_{r-s+k} \right\}, r \ge s$$

and

$$Q_n(x) = \frac{1}{\sum_{k=0}^{m} k!\binom{n-s}{k} P_{k,k+s}} \left\{ x^{n-s} - \left[\sum_{k=1}^{m} \left(\sum_{j=k}^{m} j!\binom{n-s}{j} P_{j,j-k} \right) Q_{n-s-k}(x) \right. \right. \tag{11}$$

$$\left. \left. + \sum_{k=0}^{s-1} \left(\sum_{j=0}^{m} j!\binom{n-s}{j} P_{j,j+k} \right) Q_{n-s+k}(x) \right] \right\}$$

3.0 Error estimate for the recursive form (RF)

The canonical polynomials were generated and generalized [14, 19], and this was used in error estimation of the Tau method. Adeniyi [16, 18] reported a polynomial estimate $(e_n(x))_{n+1} \cong E_{n+1}(x)$ of degree $(n+1)$ as:

$$(e_n(x))_{n+1} \cong E_{n+1}(x) = \frac{\phi_n v_m(x) T_{n-m+1}(x)}{C_{n-m+1}^{(n-m+1)}} = \frac{\phi_n v_m(x) \sum_{r=0}^{n-m+1} C_r^{(n-m+1)} x^r}{C_{n-m+1}^{(n-m+1)}} \tag{12}$$

where $v_m(x)$ is to ensure that $E_{n+1}(x)$ satisfies some or all the homogenous conditions of $e_n(x)$.

He also reported that:

$$L(E_{n+1}(x)) = \hat{H}_{n+1}(x) - H_n(x) \tag{13}$$

where

$$\hat{H}_{n+1}(x) = \sum_{i=0}^{m+s} \tau_{i+1} \sum_{r=0}^{n-m+i-2} C_r^{(n-m+i+2)} x^r \tag{14}$$

After assuming that a transformation has been made such that $a = 0, b = 1$, equation (12) becomes:

$$E_{n+1}(x)=\frac{\phi_n(x)\sum_{r=0}^{n-m+1}C_r^{(n-m+1)}x^{r+m}}{C_{n-m+1}^{(n-m+1)}} \tag{15}$$

as an approximate to the error

$$e_n(x)=y(x)-y_n(x) \tag{16}$$

in $y_n(x)$ obtained from the Tau approximation process [4, 5]. The parameter φ_n is to be determined along with $\tilde{\tau}_r$'s parameters [19] and the exact error (maximum error) is defined as:

$$\xi=a\leq^{\max} x\leq b\left\{\left|y(x)-y_n(x)\right|\right\} \tag{17}$$

4.0 Error estimate for the power series form (PSF)

The error function (16), which satisfies the perturbed error problem

$$L(e_n(x))=-H_n(x)=-\sum_{r=0}^{m+s-1}\tau_{m+s-r}T_{n-m+r+1}(x) \tag{18}$$

$$L^*e_n(\alpha)=0$$

Satisfies the perturbed error problem [17, 18, 19]

$$L(e_n(x))_{n+1}=-H_n(x)+H_{n+1}(x)=-\sum_{r=0}^{m+s-1}\tau_{m+s-r}T_{n-m+r+1}(x)+\sum_{r=0}^{m+s-1}\beta_{m+s-r}T_{n-m+r+2}(x) \tag{19}$$

$$L^*e_n(\alpha)_{n+1}=0$$

where the extra $\beta_r, r=1,2,\ldots,m+s$ parameters and φ_n are to be determined, using zeroes of chebyshev polynomial together with the given conditions. A forward elimination process is recommended for the solution of the resulting linear system. The value of φ_n is then inserted into (15) and subsequently, we get the estimate:

$$E_{n+1}(x)=a\leq^{\max} x\leq b\left|(e_n(x))_{n+1}\right|=\left|\frac{\varphi_n}{C_{n-m+1}^{n-m+1}}\right| \tag{20}$$

5.0 Numerical Examples

In this section, we applied the presented method to some selected examples. The main objective here is to solve five examples by power series using zeros of chebyshev polynomial and compare the results with method discussed in section 2. All the approximant solution are subject to degree 5.

Example 4.1

$$y''(x)+y(x)=0,\ y(0)=1,\ y'(0)=0,\ 0\leq x\leq 1$$

With analytical solution $y(x)=\cos(x)$

Example 4.2

$$Ly(x)=2(1+x)y'(x)+y(x)=0,\ y(0)=1,\ 0\leq x\leq 1$$

with analytical solution $y(x)=(1+x)^{\frac{-1}{2}}$

Example 4.3

$$Ly(x)=y'(x)-x^2y(x)=0,\ y(0)=1$$

with analytical solution $y(x)=\exp\left(\frac{1}{3}x^3\right)$

Example 4.4

$y'''(x)-y''(x)-y'(x)+y(x)=,\ y(0)=2,\ y'(0)=1, y''(0)=0, 0\leq x\leq 1$ [19]

with analytical solution $y(x)=(2-x)\exp(x)$

Example 4.5

$$y^{iv}(x)-3601y''(x)+3600y(x)=-1+1800x^2,\ y(0)=y'(0)=y''(0)=y'''(0)=1,\ 0\leq x\leq 1\ [16]$$

with analytical solution $y(x)=\frac{1}{2}\exp(-x)\left(1+2\exp(x)+\exp(2x)+x^2\exp(x)\right)$

Table 4.1: Exact Error

Method	Example 4.1 (ξ)	Example 4.2 (ξ)	Example 4.3 (ξ)	Example 4.4 (ξ)	Example 4.5 (ξ)
RF	1.16×10^{-5}	3.08×10^{-5}	2.07×10^{-4}	1.70×10^{-5}	2.77×10^{-3}
PSF	1.16×10^{-5}	1.16×10^{-5}	2.07×10^{-4}	1.70×10^{-5}	2.77×10^{-3}

Table 4.2: Error Estimate (Maximum Error)

Method	Example 4.1	Example 4.2	Example 4.3	Example 4.4	Example 4.5
RF	1.34×10^{-5}	2.90×10^{-5}	4.94×10^{-4}	4.72×10^{-4}	7.68×10^{-6}
PSF	1.20×10^{-5}	2.90×10^{-5}	4.44×10^{-4}	3.93×10^{-4}	$7.68\ 10^{-6}$

6.0 Conclusion

In this work, we have studied the power series approach using zeros of chebyshev polynomial to compute the error estimate of ODEs. The error estimate obtained with the presented method are in good agreement with the recursive form, as it is favourably compared and give accurate results.

7.0 References

[1] Adeniyi, R.B. and Edungbola, E.O. (2008), On the tau method for certain over determined first order differential equations, J. Nig. Ass. Mathematical Physics 12,399-408.

[2] Adeniyi,R.B. and Edungbola, E.O. (2007), On the recursive formulation of the tau method for class of over determined first order equations, Abacus J. Math. Assoc. Nig.,34(2B), 249-261.

[3] Fox, L. and Parker,I.B. (1968), Chebyshev Polynomials in Numerical Analysis, University Press, Oxford.

[4] Issa, K. and Adeniyi, R.B. (2013), A generalized scheme for the numerical solution of initial value problems in ordinary differential equations by the recursive formulation of Tau method, Intl. Jour. of Pure and Appl. Math.88(1), 1-13.

[5] Issa, K. and Jimoh, A. K. (2014), An Error Estimation of a numerical scheme analogous to the Tau method for initial value problems in Ordinary Differential Equations, J. Nig. Ass. Math. Phy.26, 78-81.

[6] Lanczos, C. (1938), Trigonometric interpolation of empirical and analytic functions, J. Math. and Physics, 17, 123-199.

[7] Lanczos, C. (1956), Applied analysis, Prentice-Hall, New Jersey.

[8] Ortiz, E.L. and H. Samara (1983), Numerical solution of differential eigenvalue problems with an operational approach to the Tau Method, Computing, 31, 95-103.

[9] Ortiz, E.L. and H. Samara (1981), An operational approach to the Tau Method for the numerical solution of non-linear differential equations, Computing, 27, 15-25.

[10] Sam C. N., (2004), Numerical Solution of Partial Differential Equations with the Tau Collocation Method, Master of Philosophy, City University Of Hong Kong.

[11] Crisci, M.R. and Russo, E.(1983), An extension of ortiz's recursive formulation of tau method for certain linear system of ordinary differential equation, Math Comput., 41:431-435.

[12] El-Daou, M.K. (2005),A tau method with the perturbed boundary conditions for certain ODEs, Numerical Algorithms,38: 31-45.

[13] El-Daou,M.K. and Ortiz, E.L. (2007), Weight subspaces of the tau method and orthogonal collocation, J. Math. Anal. Appl.,326: 622-631.

[14] Yisa, B.M. and Adeniyi, R.B., Generalization of canonical polynomials for over determined$m$ — th order ordinary differential equations(ODEs), IJERT, 1(6)(2012), 1-15.

[15] Yisa, B.M. and Adeniyi, R.B. (2012), On the generalization of the error and errorestimate process of Ortiz's recursive formulation of the Tau method, O.A.U. conference, Ile-Ife, Nigeria.

[16] Adeniyi, R.B. (1991), On the tau method for numerical solution of on'linary differential equations, Doctoral thesis, University of Ilorin.

[17] Adeniyi,R.B., Onumanyi, P. and Taiwo, O.A. (1990),A computational error estimate of tau method for non linear ordinary differential equations, J. Nig. Maths Soc., 21-32.

[18] Adeniyi, R. B. and Onumanyi, P. (1991); Error estimation in the numerical solution of ODE with the Tau method. Comput. and Math. Applic., 21(9), 19-27.

[19] Yisa, B.M. and Adeniyi, R.B. (2012), On the generalization of the error and error estimate process of Ortiz's recursive formulation of the Tau method, O.A.U. NMC conference Proceeding, Ile-Ife, Osun State