ResearchGate

See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/366990505

Toeplitz Determinants for a Subclass of Analytic Functions

Article · January 2023

CITATIONS
7

READS
137

3 authors, including:



Ayinla Rasheed
Kwara State University
11 PUBLICATIONS **30** CITATIONS
SEE PROFILE



Risikat Bello
Kwara State University
5 PUBLICATIONS **20** CITATIONS
SEE PROFILE

All content following this page was uploaded by Ayinla Rasheed on 10 January 2023.

The user has requested enhancement of the downloaded file.

Journal of Progressive Research in Mathematics
Vol 18,1 - 2021 — E-ISSN: 2395-0218
http://www.scitecresearch.com/journals/index.php/jprm

Toeplitz Determinants for a Subclass of Analytic Functions

AYINLA, Rasheed O BELLO, Risikat, A

Department of Statistics and Mathematical Sciences,
P.M.B 1530,
Kwara State University, Malete.
Email: rasheed.ayinla@kwasu.edu.ng

Department of Statistics and Mathematical Sciences,
P.M.B 1530,
Kwara State University, Malete.
Email: risikat.bello@kwasu.edu.ng

Received: March 07, 2021; Accepted: March 29, 2021; Published: May 3, 2021

Copyright © 2021 by author(s) and Scitech Research Organisation(SRO).
This work is licensed under the Creative Commons Attribution International License (CC BY).
http://creativecommons.org/licenses/by/4.0/

Abstract

A new subclass of analytic functions that generalizes some known subclasses of analytic functions was defined and investigated.The bounds for Toeplitz determinants of $T_2(2), T_2(3), T_3(1)$ and $T_3(2)$ were obtained.

Keywords

Analytic functions, Salagean differential operator and Toeplitz determinant.

1. Introduction

Let H denote the class of functions of the form

$$f(z) = z + a_2 z^2 + a_3 z^3 + ... \tag{1.1}$$

which are analytic in the open unit disk $E = \{z : |z| < 1\}$ and satisfy the condition $f(0) = 0$ and $f^{'}(0) = 1$. Let S denote the subclass of H consisting of univalent in E. A funtion $f(z) \in S$ is said to be starlike in the unit disk if and only if

$$Re\frac{zf'(z)}{f(z)} > 0, \qquad z \in E \tag{1.2}$$

Also, a function $f(z) \in S$ is said to be convex in the unit disk if and only if

$$Re\left(1 + \frac{zf''(z)}{f'(z)}\right) > 0, \qquad z \in E \tag{1.3}$$

Cite this paper: Ayinla, R., Bello, R. (2021). Toeplitz Determinants for a Subclass of Analytic Functions. Journal of Progressive Research in Mathematics, 18(1), 99-106. Retrieved from http://scitecresearch.com/journals/index.php/jprm/article/view/2034

Let $D^n : A \to A$ be defined by

$$D^0 f(z) = f(z)$$
$$D^1 f(z) = zf'(z)$$
$$D^n f(z) = z[D^{n-1} f(z)]'$$

which is equivalent to

$$D^n f(z) = z + \sum_{k=2}^{\infty} k^n a_k z^k, \quad (n = \{0, 1, 2, ...\}), \quad z \in E$$

D^n is the Salagean differential operator [7].
Hankel determinants play important role in several branches of mathematics such as qantum mechanics, image processing, statistics and probability, queueing networks, signal processing and time series analysis to mention a few [9]. A Toeplitz determinants is an upside down Hankel determinants, that is Hankel determinants have constant entries along the reverse diagonal while Toeplitz determinants have constant entries the diagonal.
Thomas and Halim [8] introduced the symmetric Toeplitz determinant $T_q(n)$ for analytic functions $f(z)$ of the form (1.1) defined as follows

$$T_q(n) = \begin{vmatrix} a_n & a_{n+1} & ... & a_{n+q-1} \\ a_{n+1} & a_n & ... & a_{n+q-2} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n+q-1} & a_{n+q-2} & ... & a_n \end{vmatrix} \qquad (where\ n, q = 1, 2, 3, \cdots a_1 = 1\ for f(z) \in S)$$

In particular,

$$T_2(2) = \begin{vmatrix} a_2 & a_3 \\ a_3 & a_2 \end{vmatrix}, \quad T_2(3) = \begin{vmatrix} a_3 & a_4 \\ a_4 & a_3 \end{vmatrix}, \quad T_3(1) = \begin{vmatrix} 1 & a_2 & a_3 \\ a_2 & 1 & a_2 \\ a_3 & a_2 & 1 \end{vmatrix}, T_3(2) = \begin{vmatrix} a_2 & a_3 & a_4 \\ a_3 & a_2 & a_3 \\ a_4 & a_3 & a_2 \end{vmatrix}$$

The estimates of the Toeplitz determinant $T_q(n)$ for functions in S^* and K have been studied in [8]. The Toeplitz determinant for some subclasses of analytic function was discussed in [6]. Also, estimates of the Toeplitz determinant for boundary rotation (R) was studied in [5] for small n and q.
Let P denote the class of analytic functions p in E of the form

$$p(z) = 1 + \sum_{n=1}^{\infty} p_n z^n \tag{1.4}$$

such that $Rep(z) > 0$ in E.
A function $f(z)$ of the form (1.1) analytic and univalent in E is said to be in the $R_n(\alpha, \beta)$, $\alpha \in [0, 1]$, $\beta \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$ and $n \in \mathbb{N}_0$ if it satisfies the inequality

$$Re\left\{e^{i\beta}\left(1 - e^{-2i\beta}\alpha^2 z^2\right)\frac{D^{n+1}f(z)}{z}\right\} > 0, z \in E. \tag{1.5}$$

Remark 1
(ii) For $n = 0$, $\beta = 0$ gives

$$Re\left\{\left(1 - \alpha^2 z^2\right) f'(z)\right\} > 0, \quad z \in E. \tag{1.6}$$

investigated by kanas and Lecko [2].
For $n = 0$, (1.5) gives

$$Re\left\{e^{i\beta}\left(1 - e^{-2i\beta}\alpha^2 z^2\right) f'(z)\right\} > 0, \quad z \in E. \tag{1.7}$$

studied in[3].

2 Preliminary Lemmas

We need the following lemmas to prove our results.

Let P denote the class of Caratheodory functions.
$p(z) = 1 + p_1 z + p_2 z^2 + 3_3 z^3 + ... \quad (z \in E)$
which are analytic and satisfy $p(0) = 1$ and $\Re p(z) > 0$ Let $p \in P$. Then

$$|p_k| \leq 2 \quad (k \in \mathbb{N}) \quad [1] \tag{2.1}$$

. Let $p \in P$ then

$$2p_2 = p_1^2 + x\left(4 - p_1^2\right) \tag{2.2}$$

$$4p_3 = p_1^3 + 2p_1(4-_1^2)x - p_1(4-p_1^2)x^2 + 2(4-p_1^2)(1-|x|^2)z \tag{2.3}$$

for some value of x, z, such that $|x| \leq 1$ and $|z| \leq 1$. [4]

3 Main Results

Let $f(z) \in R_n(\alpha, \beta), \alpha \in [0,1], \quad \beta \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$ and $n \in \mathbb{N}_0$. Then

$$|T_2(2)| = |a_3^2 - a_2^2| \leq \frac{4\cos^2\beta + 4\alpha^2\cos\beta}{3^{2n+2}} + \frac{\cos\beta}{2^{2n}}$$

Proof:
Let $f(z) \in R_n(\alpha, \beta)$, then there exist $p \in P$ such that

$$e^{i\beta}\left\{\left(1 - e^{-2i\beta}\alpha^2 z^2\right)\left(1 + \sum_{k=2}^{\infty} k^{n+1} a_k z^{k-1}\right)\right\} = p(z)\cos\beta + i\sin\beta \tag{3.1}$$

Equating coefficients of (3.1) give

$$a_2 = \frac{p_1 e^{-i\beta}\cos\beta}{2^{n+1}} \tag{3.2}$$

$$a_3 = \frac{p_2 e^{-i\beta}\cos\beta + \alpha^2 e^{-2i\beta}}{3^{n+1}} \tag{3.3}$$

$$a_4 = \frac{p_3 e^{-i\beta}\cos\beta + p_1\alpha^2 e^{-3i\beta}\cos\beta}{4^{n+1}} \tag{3.4}$$

Using (3.2) and (3.3) we obtain

$$|T_2(2)| = \left|a_3^2 - a_2^2\right| = \left|\frac{p_2^2 e^{-2i\beta}}{3^{2n+2}}\cos^2\beta + \frac{2p_2\alpha^2 e^{-3i\beta}}{3^{2n+2}}\cos\beta - \frac{p_1^2 e^{-2i\beta}}{2^{2n+2}}\cos^2\beta + \frac{\alpha^4 e^{-4i\beta}}{3^{2n+2}}\right| \tag{3.5}$$

Substituting for p_2 using lemma (2.2) we get

$$\begin{aligned}\left|a_3^2 - a_2^2\right| = \Bigg|&\frac{p_1^4 e^{-2i\beta}}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p_1^2 x(4-p_1^2)e^{-2i\beta}}{2\cdot 3^{2n+2}}\cos^2\beta + \frac{\alpha^2 x(4-p_1^2)e^{-3i\beta}}{3^{2n+2}}\cos\beta \\ &- \frac{p_1^2 e^{-2i\beta}}{2^{2n+2}}\cos^2\beta + \frac{x^2(4-p_1^2)^2 e^{-2i\beta}}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{\alpha^2 p_1^2 e^{-3i\beta}}{3^{2n+2}}\cos\beta + \frac{\alpha^4 e^{-4i\beta}}{3^{2n+2}}\Bigg|\end{aligned} \tag{3.6}$$

Let $p_1 = p, |p| \leq 2$ and assuming without loss of generality that $p \in [0,2]$. Also, applying the triangle inequality with $X = 4 - p_1^2$ gives

$$\begin{aligned}\left|a_3^2 - a_2^2\right| \leq &\frac{p^4}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p^2}{2^{2n+2}}\cos^2\beta + \frac{p^2|x|X}{2\cdot 3^{2n+2}}\cos^2\beta + \frac{\alpha^2|x|X}{3^{2n+2}}\cos\beta \\ &+ \frac{|x|^2X^2}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p^2\alpha^2}{2^{n+2}}\cos\beta + \frac{\alpha^4}{3^{2n+2}} = \mu(p,|x|)\end{aligned} \tag{3.7}$$

Differentiating $\mu(p,|x|)$ w.r.t $|x|$ and using elementary calculus shows that $\partial\mu(p,|x|/\partial|x|) > 0$ for $|x| \in [0,1]$ and fixed $p \in [0,2]$.This follows that $\mu(p,|x|)$ is an increasing function of $|x|$. So, $\mu(p,|x|) \leq \mu(p,1)$. Therefore,

$$\left|a_3^2 - a_2^2\right| \leq \frac{p^2}{2^{2n+2}}\cos^2\beta + \frac{1}{3^{2n+2}}\left(4\alpha^2\cos\beta + 4\cos^2\beta + \alpha^4\right) \tag{3.8}$$

Now, $\mu(|x|)$ has a maximum value

$$\frac{\cos^2\beta}{2^{2n}} + \frac{4\cos^2\beta + 4\alpha^2\cos\beta + \alpha^4}{3^{2n+2}}$$

on $[0,2]$ when p=2.
Hence,

$$\left|a_3^2 - a_2^2\right| \leq \frac{4\cos^2\beta + 4\alpha^2\cos\beta + \alpha^4}{3^{2n+2}} + \frac{\cos^2\beta}{2^{2n}}$$

Remark 2
For $n = \alpha = \beta = 0$

$$|T_2(2)| = \left|a_3^2 - a_2^2\right| \leq \frac{13}{9}$$

which is much finer than $|a_3^2 - a_2^2| \leq 5$ that was obtained in [6]. Let $f(z) \in R_n(\alpha,\beta), \alpha \in [0,1], \ \beta \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$ and $n \in \mathbb{N}_0$. Then

$$|T_2(3)| = \left|a_4^2 - a_3^2\right| \leq \frac{1+\alpha^4}{2^{4n+2}}\cos^2\beta + \frac{\alpha^2}{2^{4n+1}}\cos\beta + \frac{1}{3^{2n+2}}\left(4\cos^2\beta + 4\alpha^2\cos\beta + \alpha^4\right)$$

Proof. Using (3.3), (3.4) and lemma (2.2) to express p_2 and p_3 in terms of p_1, we get

$$\begin{aligned}
\left|a_4^2 - a_3^2\right| = \Bigg| & \frac{p_1^6 e^{-2i\beta}}{2^{4n+8}}\cos^2\beta + \frac{p_1^4(4-p_1^2)xe^{-2i\beta}}{2^{4n+6}}\cos^2\beta - \frac{p_1^4(4-p_1^2)x^2e^{-2i\beta}}{2^{4n+7}}\cos^2\beta \\
& + \frac{p_1^3(4-p_1^2)(1-|x|^2)ze^{-2i\beta}}{2^{4n+6}}\cos^2\beta + \frac{p_1^2(4-p_1^2)^2x^2e^{-2i\beta}}{2^{4n+6}}\cos^2\beta \\
& - \frac{p_1^2(4-p_1^2)x^3e^{-2i\beta}}{2^{4n+6}}\cos^2\beta + \frac{p_1^2(4-p_1^2)^2x^4e^{-2i\beta}}{2^{4n+8}}\cos^2\beta + \frac{p_1^4\alpha^2e^{-4i\beta}}{2^{4n+5}}\cos\beta \\
& + \frac{p_1(4-p_1^2)^2(1-|x|^2)xze^{-2i\beta}}{2^{4n+5}}\cos^2\beta - \frac{p_1(4-p_1^2)^2x^2(1-|x|^2)ze^{-2i\beta}}{2^{4n+6}}\cos^2\beta \\
& + \frac{(4-p_1^2)^2(1-|x|^2)^2z^2e^{-2i\beta}}{2^{4n+6}}\cos^2\beta + \frac{p_1^2(4-p_1^2)\alpha^2xe^{-4i\beta}}{2^{4n+4}}\cos\beta \\
& - \frac{p_1^2(4-p_1^2)\alpha^2x^2e^{-4i\beta}}{2^{4n+5}}\cos\beta + \frac{p_1(4-p_1^2)(1-|x|^2)\alpha^2ze^{-41\beta}}{2^{4n+4}}\cos\beta \\
& + \frac{p_1^2\alpha^4e^{-6i\beta}}{2^{4n+4}}\cos^2\beta - \frac{p_1^4e^{-2i\beta}}{2^2\cdot 3^{2n+2}}\cos^2\beta - \frac{p_1^2(4-p_1^2)xe^{-2i\beta}}{2\cdot 3^{2n+2}}\cos^2\beta \\
& - \frac{(4-p_1^2)^2x^2}{2^2\cdot 3^{2n+2}} + \frac{p_1^2\alpha^2e^{-2i\beta}}{3^{2n+2}}\cos\beta + \frac{(4-p_1^2)\alpha^2xe^{-2i\beta}}{3^{2n+2}}\cos\beta - \frac{\alpha^4e^{-4i\beta}}{3^{2n+2}}\Bigg|
\end{aligned} \tag{3.9}$$

Letting $p_1 = p, |p_1| \leq 2$, assuming without loss of generality that $p \in [0,2]$ and applying triangle inequality with $X = 4 - p_1^2$ and $Y = (1 - |x|^2)$ we obtain

$$\begin{aligned}\left|a_4^2 - a_3^2\right| \leq{}& \frac{p^6}{2^{4n+8}}\cos^2\beta + \frac{p^4|x|X}{2^{4n+6}}\cos^2\beta + \frac{p^4|x|^2X}{2^{4n+7}}\cos^2\beta + \frac{p^3XY}{2^{4n+6}}\cos^2\beta \\ &+ \frac{p^2|x|^2X^2}{2^{4n+6}}\cos^2\beta + \frac{p^2|x|^3X^2}{2^{4n+6}}\cos\beta + \frac{p^2|x|^4X^2}{2^{4n+8}} + \frac{p|x|X^2Y}{2^{4n+5}}\cos^2\beta \\ &+ \frac{p|x|^2X^2Y}{2^{4n+6}}\cos^2\beta + \frac{X^2Y^2}{2^{4n+6}}\cos^2\beta + \frac{p^4\alpha^2}{2^{4n+5}}\cos\beta + \frac{p^2|x|X\alpha^2}{2^{4n+4}}\cos\beta \\ &+ \frac{p^2|x|^2X\alpha^2}{2^{4n+5}}\cos\beta + \frac{pXY\alpha^2}{2^{4n+4}}\cos\beta + \frac{p^2\alpha^4}{2^{4n+4}}\cos^2\beta + \frac{p^4}{2^2\cdot 3^{2n+2}}\cos^2\beta \\ &+ \frac{p^2|x|X}{2\cdot 3^{2n+2}}\cos^2\beta + \frac{|x|^2X^2}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p^2\alpha^2}{3^{2n+2}}\cos\beta + \frac{\alpha^2|x|X}{3^{2n+2}}\cos\beta \\ &+ \frac{\alpha^4}{3^{2n+2}} = \phi(p,|x|) \end{aligned} \tag{3.10}$$

Differentiating $\phi(p,|x|)$ w.r.t $|x|$ and using elementary calculus shows that $\partial\phi(p,|x|/\partial|x|) > 0$ for $|x| \in [0,1]$ and fixed $p \in [0,2]$.This follows that $\phi(p,|x|)$ is an increasing function of $|x|$. So, $\phi(p,|x|) \leq \phi(p,1)$.

$$\begin{aligned}\left|a_4^2 - a_3^2\right| \leq{}& \frac{p^6}{2^{4n+8}}\cos^2\beta + \frac{3p^4(4-p^2)}{2^{4n+7}}\cos^2\beta + \frac{5p^2(4-p^2)^2}{2^{4n+8}}\cos^2\beta + \frac{p^4\alpha^2}{2^{4n+5}}\cos\beta \\ &+ \frac{3p^2(4-p^2)\alpha^2}{2^{4n+5}}\cos\beta + \frac{p^2\alpha^4}{2^{4n+4}}\cos^2\beta + \frac{p^4}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p^2(4-p^2)}{2\cdot 3^{2n+2}}\cos^2\beta \\ &+ \frac{(4-p^2)^2}{2^2\cdot 3^{2n+2}}\cos^2\beta + \frac{p^2\alpha^2}{3^{2n+2}}\cos\beta + \frac{(4-p^2)\alpha^2}{3^{2n+2}}\cos\beta + \frac{\alpha^4}{3^{2n+2}} \end{aligned} \tag{3.11}$$

(3.11) has maximum value

$$\frac{1+\alpha^4}{2^{4n+2}}\cos^2\beta + \frac{\alpha^2}{2^{4n+1}}\cos\beta + \frac{1}{3^{2n+2}}\left(4\cos^2\beta + 4\alpha^2\cos\beta + \alpha^4\right)$$

on $[0,,2]$ when $p = 2$.
Remark 3: When $n = \alpha = \beta = 0$,

$$|T_2(3)| = \left|a_4^2 - a_3^2\right| \leq \frac{25}{36}$$

. which is much finer than $|a_4^2 - a_3^2| \leq 7$ that was obtained in [6] for the class of function with real positive part. Let $f(z) \in R_n(\alpha,\beta), \alpha \in [0,1], \quad \beta \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$ and $n \in \mathbb{N}_0$. Then

$$\begin{aligned}|T_3(2)| = \left|(a_2 - a_4)(a_2^2 - 2a_3^2 + a_2a_4)\right| \\ \leq \left[\frac{1+\alpha^2}{2^{2n+1}}\cos\beta + \frac{1}{2^n}\cos\beta\right]\left[\frac{1}{2^{2n}}\cos^2\beta + \frac{(\alpha^2-1)}{2^{3n+1}}\cos^2\beta\right. \\ \left. + \frac{8}{3^{2n+2}}\cos^2\beta - \frac{8\alpha^2}{3^{2n+2}}\cos\beta + \frac{2\alpha^4}{3^{2n+2}}\right]\end{aligned}$$

Proof.

$$|T_3(2)| = \left|(a_2 - a_4)(a_2^2 - 2a_3^2 + a_2a_4)\right| \leq |a_2 - a_4|\left|a_2^2 - 2a_3^2 + a_2a_4\right|$$

Now, using (3.2), (3.4) and lemma (2.2)

$$\begin{aligned}|a_2 - a_4| = \left|\frac{p_1e^{-i\beta}}{2^{n+1}}\cos\beta - \frac{p_1^3e^{-i\beta}}{2^{2n+4}}\cos\beta - \frac{p_1(4-p_1^2)xe^{-i\beta}}{2^{2n+3}}\cos\beta\right. \\ \left. + \frac{p_1(4-p_1^2)x^2e^{-i\beta}}{2^{2n+4}}\cos\beta - \frac{(4-p_1^2)(1-|x|^2)ze^{-i\beta}}{2^{2n+3}}\cos\beta - \frac{p_1\alpha^2e^{-3i\beta}}{2^{2n+2}}\cos\beta\right|\end{aligned}$$

Using triangle inequality with $p_1 = p$

$$|a_2 - a_4| \leq \frac{p^3}{2^{2n+4}}\cos\beta + \frac{p}{2^{n+1}}\cos\beta + \frac{p|x|X}{2^{2n+3}}\cos\beta + \frac{p|x|^2X}{2^{2n+4}}\cos\beta + \frac{XY}{2^{2n+3}}\cos\beta + \frac{p\alpha^2}{2^{2n+2}}\cos\beta$$

Using the same techniques as in theorem (3.1) and (3.2)

$$|a_2 - a_4| \leq \frac{1+\alpha^2}{2^{2n+1}}\cos\beta + \frac{1}{2^n}\cos\beta$$

From (3.2),(3.3) and (3.4) we get

$$\left|(a_2^2 - 2a_3^2 + a_2a_4)\right| = \left|\frac{p_1^2e^{-2i\beta}}{2^{2n+2}}\cos^2\beta - \frac{2p_2^2e^{-2i\beta}}{3^{2n+2}}\cos^2\beta - \frac{4p_2\alpha^2e^{-3i\beta}}{3^{2n+2}}\cos\beta - \frac{2\alpha^4e^{-4i\beta}}{3^{2n+2}} + \frac{p_1p_3e^{-2i\beta}}{2^{3n+3}}\cos^2\beta + \frac{p_1^2\alpha^2e^{-4i\beta}}{2^{3n+3}}\cos^2\beta\right|$$

Letting $p_1 = p \in [0,2]$ and applying triangle inequality and simplifying, we obtain

$$\begin{aligned}\left|(a_2^2 - 2a_3^2 + a_2a_4)\right| &\leq \left[\frac{\cos^2\beta}{2^{2n+2}} - \frac{-2\alpha^2\cos\beta}{3^{2n+2}} + \frac{\alpha^2\cos^2\beta}{2^{3n+3}}\right]p^2 + \left[\frac{\cos^2\beta}{2\cdot 3^{2n+2}} - \frac{\cos^2\beta}{2^{3n+5}}\right]p^4 \\ &+ \left[\frac{\cos^2\beta}{3^{2n+2}} - \frac{\cos^2\beta}{2^{3n+4}}\right](4-p^2)|x|p^2 + \frac{2\alpha^2(4-p^2)|x|}{3^{2n+2}}\cos\beta \\ &+ \frac{p^2(4-p^2)|x|^2}{2^{3n+5}}\cos^2\beta + \frac{(4-p^2)^2|x|^2}{2\cdot 3^{2n+2}}\cos^2\beta \\ &+ \frac{p(4-p^2)(1-|x|^2)}{2^{3n+4}}\cos^2\beta + \frac{2\alpha^4}{3^{2n+2}} = \gamma(p,|x|)\end{aligned}$$

Now, we need to obtain the maximum value of $\gamma(p,|x|)$ on the closed region $[0,2]X[0,1]$. Let us assume that there is a maximum at an interior point $(p_0,|x|)$ of $[0,2]X[0,1]$. Then, differentiating $\gamma(p_0,|x|)$ w.r.t $|x|$ we get

$$\begin{aligned}\frac{\partial\gamma}{\partial|x|} &= \left[\frac{\cos^2\beta}{3^{2n+2}} - \frac{\cos^2\beta}{2^{3n+4}}\right](4-p^2)p^2 + \frac{2\alpha^2(4-p^2)}{3^{2n+2}} + \frac{p^2(4-p^2)|x|}{2^{3n+4}}\cos^2\beta \\ &+ \frac{(4-p^2)^2|x|}{3^{2n+2}}\cos^2\beta - \frac{p(4-p^2)|x|}{2^{3n+3}} \qquad (3.12)\end{aligned}$$

equating (3.12) to zero imply that $p = 2$ which is a contradiction. Thus, we need to consider only the endpoints of $[0,2]X[0,1]$ in other to obtain the maximum of $\gamma(p,|x|)$.
When $p = 0$,

$$\gamma(0,|x|) = \frac{8\alpha^2|x|}{3^{2n+2}}\cos\beta + \frac{16|x|^2}{2\cdot 3^{2n+2}}\cos^2\beta + \frac{2\alpha^4}{3^{2n+2}} \leq \frac{8\alpha^2}{3^{2n+2}}\cos\beta + \frac{8}{3^{2n+2}}\cos^2\beta + \frac{2\alpha^4}{3^{2n+2}}$$

When $p = 2$,

$$\gamma(2,|x|) = \frac{4}{2^{2n+2}}\cos^2\beta - \frac{8\alpha^2}{3^{2n+2}}\cos\beta + \frac{4\alpha^2}{2^{3n+3}}\cos^2\beta + \frac{8}{3^{2n+2}}\cos^2\beta - \frac{1}{2^{3n+1}}\cos^2\beta + \frac{2\alpha^4}{3^{2n+2}}$$

When $|x| = 0$,

$$\begin{aligned}\gamma(p,0) &= \left[\frac{\cos^2\beta}{2^{2n+2}} - \frac{2\alpha^2\cos\beta}{3^{2n+2}} + \frac{\alpha^2\cos^2\beta}{2^{3n+3}}\right]p^2 + \left[\frac{\cos^2\beta}{2\cdot 3^{2n+2}} - \frac{\cos^2\beta}{2^{3n+5}}\right]p^4 \\ &+ \frac{p(4-p^2)}{2^{3n+4}}\cos^2\beta + \frac{2\alpha^4}{3^{2n+2}}\end{aligned}$$

which has a maximum value

$$\frac{1}{2^{2n}}\cos^2\beta-\frac{8\alpha^2}{3^{2n+2}}\cos\beta+\frac{\alpha^2}{2^{3n+1}}\cos^2\beta-\frac{1}{2^{3n+1}}\cos^2\beta+\frac{2\alpha^4}{3^{2n+2}}$$

on $[0,2]$. Also, when $|x|=1$

$$\begin{aligned}\gamma(p,1)=&\left[\frac{\cos^2\beta}{2^{2n+2}}-\frac{2\alpha^2\cos\beta}{3^{2n+2}}+\frac{\alpha^2\cos^2\beta}{2^{3n+3}}\right]p^2+\left[\frac{\cos^2\beta}{2\cdot 3^{2n+2}}-\frac{\cos^2\beta}{2^{3n+5}}\right]p^4\\&+\left[\frac{\cos^2\beta}{3^{2n+2}}-\frac{\cos^2\beta}{2^{3n+4}}\right]p^2(4-p^2)+\frac{2\alpha^2(4-p^2)}{3^{2n+2}}\cos\beta+\frac{p^2(4-p^2)}{2^{3n+5}}\cos^2\beta\\&+\frac{(4-p^2)^2}{2\cdot 3^{2n+2}}\cos^2\beta+\frac{2\alpha^4}{3^{2n+2}}\end{aligned}$$

which has a maximum value

$$\frac{1}{2^{2n}}\cos^2\beta-\frac{8\alpha^2}{3^{2n+2}}\cos\beta+\frac{\alpha^2}{2^{3n+1}}+\frac{8}{3^{2n+2}}\cos^2\beta-\frac{1}{2^{3n+1}}\cos^2\beta+\frac{2\alpha^4}{3^{2n+2}}$$

on $[0,2]$. Hence,

$$\begin{aligned}|T_3(2)|=&\left|(a_2-a_4)(a_2^2-2a_3^2+a_2a_4)\right|\\&\leq\left[\frac{1+\alpha^2}{2^{2n+1}}\cos\beta+\frac{1}{2^n}\cos\beta\right]\left[\frac{1}{2^{2n}}\cos^2\beta+\frac{(\alpha^2-1)}{2^{3n+1}}\cos^2\beta\right.\\&\qquad\left.+\frac{8}{3^{2n+2}}\cos^2\beta-\frac{8\alpha^2}{3^{2n+2}}\cos\beta+\frac{2\alpha^4}{3^{2n+2}}\right]\end{aligned}$$

Remark 4 For $n=\alpha=\beta=0$,

$$|T_3(2)|=\left|(a_2-a_4)(a_2^2-2a_3^2+a_2a_4)\right|\leq\frac{25}{12}$$

which is also finer than $\left|(a_2-a_4)(a_2^2-2a_3^2+a_2a_4)\right|\leq 12$ obtained in [6] for functions with real positive part. Let $f(z)\in R_n(\alpha,\beta),\alpha\in[0,1],\ \ \beta\in\left(\frac{-\pi}{2},\frac{\pi}{2}\right)$ and $n\in\mathbb{N}_0$. Then

$$|T_3(1)|=\left|1+2a_2^2(a_3-1)-a_3^2\right|\leq 1+\frac{(4-2\alpha^2)}{2^{2n}\cdot 3^{n+1}}\cos^2\beta+\frac{2}{2^{2n}}\cos^2\beta+\frac{(4\alpha^2\cos\beta-4\cos^2\beta+\alpha^4)}{3^{2n+2}}$$

Proof. By using (3.2), (3.3) , lemma (2.2) and simplifying, we get

$$\begin{aligned}T_3(1)=\left|1+2a_2^2(a_3-1)-a_3^2\right|=&\left|1+\left[\frac{e^{-3i\beta}\cos^2\beta}{2^{2n+2}\cdot 3^{n+1}}-\frac{e^{-2i\beta}\cos^2\beta}{2^2\cdot 3^{2n+2}}\right]p_1^4\right.\\&-\left[\frac{e^{-2i\beta}\cos^2\beta}{2^{2n+1}}+\frac{\alpha^2e^{-i\beta}\cos\beta}{3^{2n+2}}-\frac{\alpha^2e^{-4i\beta}\cos^2\beta}{2^{2n+1}\cdot 3^{n+1}}\right]p_1^2\\&\left.-\left[\frac{e^{-i\beta}\cos^2\beta}{2\cdot 3^{2n+2}}-\frac{e^{-3i\beta}\cos^2\beta}{2^{2n+2}\cdot 3^{n+1}}\right]xXp_1^2+\frac{e^{-3i\beta}\cos^2\beta}{2^2\cdot 3^{2n+2}}x^2X^2-\frac{\alpha^2e^{-i\beta}\cos\beta}{3^{2n+2}}xX-\frac{\alpha^4e^{-4i\beta}}{3^{2n+2}}\right|\end{aligned}$$

Assuming without loss of generality that $p_1=p\in[0,2]$ and $X=(4-p^2)$. Also, using the triangle inequality and the fact that $|x|=1$, we get

$$\begin{aligned}|T_3(1)|\leq&1+\left[\frac{\cos^2\beta}{2^{2n+2}\cdot 3^{n+1}}-\frac{\cos^2\beta}{2^2\cdot 3^{2n+2}}\right]p^4+\left[\frac{\cos^2\beta}{2^{2n+1}}+\frac{\alpha^2\cos\beta}{3^{2n+2}}-\frac{\alpha^2\cos^2\beta}{2^{2n+1}\cdot 3^{n+1}}\right]p^2\\&+\left[\frac{\cos^2\beta}{2\cdot 3^{n+2}}-\frac{1}{2^{2n+2}\cdot 3^{n+1}}\right](4-p^2)p^2+\frac{\cos^2\beta(4-p^2)^2}{2^2\cdot 3^{2n+2}}\\&+\frac{\alpha^2\cos\beta}{3^{2n+2}}(4-p^2)+\frac{\alpha^4}{3^{2n+2}}\end{aligned}\tag{3.13}$$

The RHS of (3.13) has a maximum at $p = 2$, therefore,

$$|T_3(1)| \leq 1 + \frac{(4 - 2\alpha^2)}{2^{2n} \cdot 3^{n+1}} \cos^2 \beta + \frac{2}{2^{2n}} \cos^2 \beta + \frac{(4\alpha^2 \cos \beta - 4 \cos^2 \beta + \alpha^4)}{3^{2n+2}}$$

Remark 5 For $n = \alpha = \beta = 0$

$$T_3(1) = \left|1 + 2a_2^2(a_3 - 1) - a_3^2\right| \leq \frac{35}{9}$$

which is finer than the result in [6].

4 Competing Interests

The authors declare that there are no competing interests regarding the publication of this paper.

5 Acknowledgements

The authors wish to thank the referees for their valuable suggestions that lead to improvement of the quality of the work in this paper.

References

[1] Duren, P.L. (1983). Univalent functions. Graduate texts in mathematics. Springer-Verlag. New York Inc. New York.

[2] Lecko, A and Kanas, S. (1990). On the Fekete-Szego problem and the domain of convexity for a certain class of univalent functions. Folia Scientiarum Universitatis Technical Resolviensis, 73, 49-56.

[3] Lecko, A. (1993). Some generalizations of analytic condition for class of convex in a given direction. Folia Scientiarum Universitatis Technical Resolviensis, 121(14), 23-24.

[4] Libera, R.J and Zlotkiewicz, E.J. (1983). Coefficient bounds for the inverse of a function with derivative. Proceedings of American Mathematical Society, 87(2), 251-257.

[5] Radhika, V. Sivasubramanian, S. Murugusundaramooorthy, G and Jahangiri, Jay M. (2016). Toeplitz matrices whose elements are the coefficients of functions with bounded boundary rotation. Journal of Complex Analysis, http//dx.doi.org/10.1155/2016/4960704, 1-4.

[6] Ramachandran, C and Kavitha, D. (2017). Toeplitz determinant for some subclasses of analytic functions. Global Journal of Pure and Applied Mathematics, 13 (2), 785-793.

[7] Salagean, S. (1983). Subclasses of univalent functions. Lecture notes in Mathematics Springer,Berlin, 1013, 362-372.

[8] Thomas, D.K. and Halim, S.A. (2016). Toeplitz matrices whose elements are the coefficients of starlike and close-to-convex functions. Bulleting of the Malaysian Mathematical Sciences Society, 193-217

[9] Ye, K. and Lim, L.K (2016). Every matrix is a product of Toeplitz matrices.Foundations of Computational Mathematics,16(3), 577-598.

View publication stats